

A Review of the Effects of Macroalgae in Shrimp Feeds and in Co-Culture

L. Elizabeth Cruz-Suárez*, Mireya Tapia-Salazar, Martha G. Nieto-López,
Denis Ricque-Marie

Programa Maricultura, Universidad Autónoma de Nuevo León, Cd. Universitaria
F-56, San Nicolás de los Garza, Nuevo León 66450, México.

E-mail: lucruz@fcb.uanl.mx

Abstract

Most nutritional studies with seaweed meals or seaweed extracts have investigated low dietary inclusion rates (less than 10%) to establish their possible usefulness as functional (binder effect), nutritional, and nutraceutical (health protective effect) supplements in shrimp feeds. The optimum inclusion level varies depending on algae or consumer species. In many instances, the inclusion of algae in feed formulations has resulted in improved pellet quality (water stability, water holding capacity and texture), higher feed intake, improved feed efficiency, better growth performance and higher animal product quality (higher pigmentation, lower cholesterol content). Additionally, macroalgae contain some active compounds that can improve animal resistance against bacterial and virus diseases. Some seaweed species can be co-cultivated with shrimps, resulting in a sustainable alternative to reduce the need for artificial feed. In this paper the studies on the effects of seaweed inclusion in shrimp feeds or seaweed co-culture with shrimp will be reviewed.

Keywords: Shrimp, seaweed, macroalgae, feed, co-culture

Introduction

Several macro algae (*Ulva*, *Undaria*, *Ascophyllum*, *Porphyra*, *Sargassum*, *Polycavernosa*, *Gracilaria* and *Laminaria*) are widely used in fish diets and there are a number of studies into their effects that have been reviewed by Nakagawa and Montgomery (2007). In this paper we will focus on studies on shrimp; however, we provide occasional reference to some work on other aquatic species.

Some macroalgae that have been evaluated in shrimp feeds are *Macrocystis pyrifera*, *Ascophyllum nodosum*, *Kappaphycus alvarezii*, *Sargassum sp*, *Gracilaria heteroclada*, *Gracilaria cervicornis*, *Caulerpa sertularioides*, *Ulva clathrata*, *Enteromorpha sp.*, *Hypnea cervicornis*, *Cryptonemia crenulata* and *Chnoospora minima*. The nutritional studies in shrimps that have investigated the effects of seaweed meal are summarized in Table 1.

Table 1. Studies on the use of seaweed meal in shrimp feeds

Shrimp species	Title	References
<i>P. monodon</i>	Use of seaweed meals from <i>Kappaphycus alvarezii</i> and <i>Gracilaria heteroclada</i> as binders in diets of juvenile shrimp <i>Penaeus monodon</i>	Peñaflorida and Golez, 1996
<i>P. monodon</i>	The potential of <i>Gracilaria spp.</i> meal for supplementation of diets for juvenile <i>Penaeus monodon</i> Fabricius	Briggs and Funge-Smith, 1996
<i>P. californiensis</i>	Efecto de la macrolaga <i>Caulerpa sertularioides</i> en el desarrollo del camarón café	Porchas Cornejo <i>et al.</i> , 1999
<i>L. vannamei</i>	Uso de harina de kelp <i>Macrocystis pyrifera</i> en alimentos para camarón	Cruz-Suárez <i>et al.</i> , 2000
	Water stability and texture of shrimp pelleted feeds formulated with natural and synthetic binders	Cruz-Suárez <i>et al.</i> , 2002b
<i>L. vannamei</i>	Inclusión de harina de kelp (<i>Macrocystis pyrifera</i>) en alimentos balanceados para el camarón	Rivera, <i>et al.</i> , 2002
<i>L. vannamei</i>	Uso de alga <i>Sargassum sp.</i> en la elaboración de dietas para camarón y su impacto en los parámetros productivos	Cruz Suárez <i>et al.</i> , 2003
<i>L. vannamei</i>	<i>Sargassum spp.</i> como fuente potencial de alimento para camarón	Casas- Valdéz <i>et al.</i> , 2002
<i>F. californiensis</i>	Efecto del alga marina <i>Sargassum spp.</i> sobre las variables productivas y la concentración de colesterol en el camarón café	Casas-Valdéz <i>et al.</i> , 2006
<i>L. vannamei</i>	Efecto de la inclusión de alginato y harina de algas <i>Sargassum sp</i> y <i>Macrocistys pyrifera</i> sobre la estabilidad en agua, digestibilidad del alimento y sobre el crecimiento del camarón blanco	Suárez-García, 2006
<i>L. vannamei</i>	Inclusión de harina de Kelp (<i>Macrocystis pyrifera</i>) en alimentos balanceados para camarón	Marinho-Soriano <i>et al.</i> , 2007
<i>L. vannamei</i>	Comparison of <i>Ulva clathrata</i> and the kelps <i>Macrocystis pyrifera</i> and <i>Ascophyllum nodosum</i> as ingredients in shrimp feeds	Cruz- Suárez <i>et al.</i> , 2008b
<i>L. vannamei</i>	Seaweed meal as a protein source for the white shrimp <i>Litopenaeus vannamei</i>	Da Silva and Barbosa, 2008
	Harina de Kelp	Cruz-Suárez, <i>et al.</i> , 2007b
<i>L. vannamei</i>	Uso de harinas de sargaso (<i>Sargassum spp.</i>) y kelp (<i>Macrocystis Pyrifera</i>) en alimentos balanceados para el camarón <i>Litopenaeus vannamei</i> efectos sobre el crecimiento y la digestibilidad <i>in vivo</i>	Gutiérrez –Leyva, 2006

Chemical composition of seaweeds

The chemical composition of macroalgae varies with species, physiological status and environmental conditions; however, in general, the macroalgae are rich in nonstarch polysaccharides, vitamins and minerals (Mabeau and Fleurence 1993; Wong and Cheung 2000). In most cases, the seaweeds are used in human or animal foods for their mineral contents or for the functional properties of their polysaccharides. Seaweeds are rarely promoted for the nutritional value of their proteins (Fleurence, 1999). The protein content of seaweed differs according to species and the seasonal period. Generally, the protein fraction of brown seaweeds is low (3 to 15% of the dry weight) compared with that of the green or red seaweeds (10 to 47% of the dry weight).

The content of crude protein, crude lipid, ash and fiber in *Macrocystis pyrifera* seaweed meals range from 5 to 14, from .5 to 2, from 31 to 45 and from 5 to 9 % respectively (Cruz-Suárez *et al.*, 2000; Rodríguez-Montesinos and Hernández-Carmona, 1991; Castro-Gonzalez *et al.*, 1994; Castro-Gonzalez *et al.*, 1991; Cruz-Suárez *et al.*, 2008b), while, for *Ascophyllum* seaweed meals, these values varied from 5 to 10, from 2 to 7, from 15 to 21 and ≈ 8 % respectively (Sharp, 1987; Cruz-Suárez *et al.*, 2008b) and for green seaweed meals from 7 to 29, from 0.5 to 4, from 13 to 36, and from 3 to 6 % respectively (Hashim and Mat-Saat, 1992; Wahbeh, 1997; Ventura and Castañón, 1998; Wong and Cheung, 2000, Wong and Cheung, 2001a; Aguilera-Morales *et al.*, 2005; Marsham *et al.*, 2007; Cruz-Suárez *et al.*, 2008b).

The amino acid composition of seaweeds has been frequently studied. For most seaweeds, aspartic and glutamic acids constitute together a large part of the amino acid fraction. In brown seaweeds, these two amino acids can represent between 22 and 44% of the total amino acids; in the green seaweeds up to 26 and 32% and in the red seaweed species 14 to 19% of the total amino acids (Fleurence, 1999). The fatty acid and pigment composition of seaweeds also differ between groups; brown and red seaweeds are a better potential sources of EPA and DHA than green ones (Ackman, 1981).

Brown seaweeds generally contain more vitamin C than their red and green counterparts. A summary of the typical whole composition of brown algae most commonly used in aquafeeds has been published by Cruz-Suárez *et al.* (2007b, 2008b).

The chemical composition of macroalgae reported in shrimp feed studies are presented in Table 2. Major nutrient (protein, carbohydrate and lipid) limitations may explain why the exclusive use of macroalgal feed (i.e. *G. cervicornis* meal) was unable to support growth and survival of white shrimp (Marinho-Soriano *et al.*, 2007).

Table 2 Proximal composition of macroalgae reported in shrimp feed studies.

Proximate composition (%)							
	Moisture	Crude protein	Crude fat	Crude fiber	NFE	Ash	
<i>Kappaphycus alvarezii</i>	10.1	3.2	0.6	5.9	72.3	18.1	Peñaflorida <i>et al.</i> , 1996
<i>Gracilaria heteroclada</i>	9.3	17.3	1.8	4.6	54.6	21.7	
<i>Macrocystis pyrifera</i>	7.4	6.1	0.7	10.5	44.2	31.1	Cruz-Suárez <i>et al.</i> , 2000
<i>Sargassum sp</i>	9.7	6.3	0.4	5.6	46.0	32.0	Casas-Valdéz <i>et al.</i> , 2002
<i>Sargassum sp.</i>	8.7	6.1	0.3	6.8	52.0	34.0	Casas-Valdéz <i>et al.</i> , 2006
<i>Gracilaria cervicornis</i>		22.9	0.5		63.1		Marinho-Soriano <i>et al.</i> , 2007
<i>Ascophyllum nodosum</i>	14.6	7.9	2.7	3.5	50.1	21.2	Cruz-Suárez <i>et al.</i> , 2008b
<i>Ulva clathrata</i>	14.2	23.4	1.0	4.6	40.8	16.0	
<i>Macrocystis pyrifera</i>	11.2	7.7	2.0	9.3	38.9	31.0	
<i>Cryptonemia crenulata</i>	18.7	21.5	1.1		44.9	13.7	Da Silva and Barbosa, 2008
<i>Hypnea cervicornis</i>	24.3	19.6	1.0		41.5	13.7	
<i>Caulerpa sertularioides fresh</i>	91.1	2.4	0.5			2.2	Porchas <i>et al.</i> , 1999
<i>Ulva clathrata fresh</i>	90.0	2.2	0.2	0.6	3.5	4.5	Cruz-Suárez <i>et al.</i> , 2008a

Pellet quality

Several studies reported that seaweed meals can be binder for aquatic feeds (Table 3). The inclusion of algae in feed formulations has resulted in improved pellet quality (hydrostability, water holding capacity and texture), resulting in higher feed intake and improved feed efficiency. The optimum inclusion level varies depending on algae or consumer species.

Briggs and Funge-Smith (1996) investigated the effects of substituting wheat flour and soybean meal with various inclusion levels (0 to 30%) of the red seaweed *Gracilaria sp.* meal on the shrimp diet stability. Inclusion at up to 10% had no significant effects on diet water stability (after 12 h), compared with the control diet lacking seaweed. Diets containing 0-15% *Gracilaria* meal remained >88% water stable after 12 h. The 30% inclusion levels of *Gracilaria* meal resulted in a significant deterioration in diet water stability (86% after 12 h). Peñaflores and Golez (1996) reported diet stabilities about 93-94% and 88% in shrimp diets supplemented with 5 to 10% *K. alvarezii* or *G. heteroclada* meals after one and four hour immersion in seawater. Cruz-Suárez *et al.* (2000) observed that the inclusion of 3 % of kelp meal resulted in better or similar shrimp pellet feed stability than for diets supplemented with a synthetic binder when feed were processed with a steam pelletizer. Marinho-Soriano *et al.* (2007) evaluated a feed made entirely of the rodophyta *Gracilaria cervicornis*, or a commercial shrimp feed; the stability in water (%) of the diets varied from approximately 82.6 % after 1h to 82.0% after 4 h of immersion in aerated seawater for *Gracilaria* pellets, and from approximately 91% after 1h to 89 % after 4h for commercial pellets. Recently Cruz Suárez *et al.* (2008b) showed that *Ulva* meal presented better binding properties than *Ascophyllum* and *Macrocystis* meals when included at 3.3% in shrimp diets.

The inclusion of macroalgae meal in feeds also affects pellet water absorption capacity (Table3). Kelp meal tends to increase the pellet water absorption, while artificial binders (Cruz-Suárez *et al.*, 2000; Cerecer-Cota *et al.*, 2005) significantly reduce this parameter. Seaweed's water absorption and gelling or binding capacity is modulated by the type and quantity of polysaccharides present (Percival, 1968; Sharp, 1987; Rodríguez-Montesinos and Hernández-Carmona, 1991; Ray and Lahaye, 1995a,b; Suzuki *et al.*, 1996; Kuda *et al.*, 1997; Paradossi *et*

al., 1999; Jimenez-Escrig and Sanchez-Muñiz, 2000; Wong and Cheung, 2001b; Marais and Joseleau, 2001; Obluchinskaia *et al.*, 2002; McHugh, 2003). When pure alginate is used as binder, the pellet holding water capacity is higher than when whole kelp (*Macrocystis* or *Sargassum*) meal is used (Cruz-Suárez *et al.*, 2000; Suárez-García, 2006). Higher pellet water absorption was found for diets with *Ulva* than for diets with *Macrocystis* and *Ascophyllum* (132 vs 112%) (Cruz-Suárez *et al.*, 2008b). The inclusion of 3.5% kelp meal in shrimp feeds produces a soft pellet texture, after immersion in water, increasing the feed intake (Cerecer-Cota, 2005).

T

able 3. Effect of macroalgae meal inclusion on shrimp pellet quality.

Algae meal	Inclusion level %	feed technology	dietary protein %	ellet water stability %1h	pellet water absorption % 1h	Ref.	
	Control starch	mincer		94		1	
<i>Kappaphycus alvarezii</i>	3,5,7,10		38-40	94			
<i>Gracilaria heteroclada</i>	3,5,7,10			93			
	Control wheat	mincer		94			
<i>Kappaphycus alvarezii</i>	5,10			94			
<i>Gracilaria heteroclada</i>	5, 10			93			
<i>Gracilaria spp.</i>	5 to 30			(0 to 15%) > 88 (30%) 86		2	
<i>Macrocystis pyrifer</i> Chile	Control wheat gluten	mincer	35	82.2		3	
	4			82			
	8			77.7			
<i>Macrocystis pyrifer</i> Mexico	Control alginate	mincer	30	91.8	180		
	2			88.8	130		
	4			87.4	150		
<i>Macrocystis pyrifer</i> Mexico	Control synthetic binder	steam pelleted	40	91.5	70		
	3.2			25	94.3		104
	3.2			25	95.6		-
<i>Macrocystis pyrifer</i> Mexico	Control Synt. binder	steam pelleted	30,30,35,40	97,98,98,96	71,65,65,66		
	3		30,30,35,40	97,98,97,96	117,102,121,130		
	5		30,30,35,40	95,97,96,95	130,135,145,137		
	3+synt		40	96	93		
	alginate	mincer	30	91.7a	191 d	4	
<i>Sargassum sp.</i>	2			88.1b	129 ab		
	4			80.9c	139 b		
<i>Macrocystis pyrifer</i>	4			87.1b	153 c		
	control			87.7b	117 a		
<i>Gracilaria cervicornis</i>	0	pelleted mincer	35	91		5	
	100		23	82.6			
<i>Ulva clathrata</i>	3.3	mincer	34	96.3	132	6	
<i>Ascophyllum nodosum</i>				95.6	112		
<i>Macrocystis pyrifer</i>				94	112		

1) Peñaflorida *et al.*, 1996; 2) Briggs and Funge, 1996; 3) Cruz- Suárez *et al.*, 2000; 4) Suárez-García, 2006; 5) Marinho-Soriano *et al.*, 2007; 6) Cruz- Suárez *et al.*, 2008b

Shrimp performance

In Table 4 are presented the studies where the effects of macroalgae meal dietary inclusion have been considered in terms of shrimp performance. In the majority of studies the inclusion of macroalgae meal improves growth, feed intake and protein efficiency at low inclusion levels (<10%).

Peñaflorida and Golez (1996) observed the best weight gain in small shrimp *P. monodon* (200 mg) fed diet including 5% *Kappaphycus alvarezii* meal and the lowest growth with the supplementation of 3% *Gracilaria heteroclada* meal; however, in a second feeding trial (500 mg initial weight), differences were not significant. Briggs and Funge-Smith (1996) found that shrimp-SGRs were very high with diets containing 0 to 15% *Gracilaria*, but lower with a 30% inclusion level; this negative effect was attributed to the high ash content, low protein content, or high level of soluble fiber present in the experimental diet with a high seaweed inclusion level. Rivera *et al.* (2002) also reported positive effects on growth of small shrimp *L. vannamei* when 10% of *Macrocystis* meal dissolved in water was used to coat pelletized diet, and poorer growth with higher (15 and 20%) seaweed levels.

On the other hand, although a significant proportion (50%) of the daily ration offered to *L. vannamei* was replaced by *Gracilaria* seaweed in the Marinho-Soriano *et al.* study (2007), the growth performance achieved (SGR =4.7%) was not significantly different with the values reported for the shrimp fed a commercial diet (5.1%).

Cruz-Suárez *et al.* (2000) found a significant increase in growth rate (53-68%) in white shrimp *L. vannamei* juveniles (450 mg) fed diets containing 2 or 4% of Mexican kelp meal (*Macrocystis pyrifera*) compared to a control diet. Nevertheless, when Chilean kelp meal (*Macrocystis pyrifera*) was tested (4 and 8% inclusion levels) in *L. vannamei* (643 mg), a slight increment in weight gain, but not significant, was observed. Suárez-García (2006) concluded that the inclusion of *Sargassum sp* meal (2-4% inclusion levels) or kelp meal *M. pyrifera* (4% inclusion level) produced growth similar to the control diet or a diet containing 3% of pure alginate as a binder. The same result was obtained by Casas-Valdez *et al.* (2006) with 4% *Sargassum* meal diets fed to shrimp *Farfantepenaeus californiensis*. In contrast, Gutiérrez-Leyva (2006) found an

improvement of SGF of shrimp *L. vannamei* fed diets with increasing levels (1, 4, 7, 10%) of *Macrocystis* or *Sargassum* meals, with 10% inclusion level giving the best growth rate. Feeding shrimp a diet supplemented with 3.3% *Ulva* meal resulted in better growth than feeding them diets supplemented with *Ascophyllum* and *Macrocystis* meals, however differences were not significant (Cruz-Suárez *et al.*, 2008b). The rhodophytes *Hypnea cervicornis* and *Cryptonemia crenulata* used as a source of protein (seaweed powder) in shrimp diets at different percentages: 39%; 26%, 13%, and 0% (control diet) exhibited no significant differences between treatments ($P>0.05$) in terms of final biomass, biomass gain and specific growth rate (Da Silva and Barbosa, 2008).

The active compound of macroalgae responsible for growth improvement has not been clearly defined and the benefit has been attributed to their vitamin and mineral content, lipid mobilization and improved absorption and assimilation efficiency ratios. The seasonal and species variations of this active compound and difference in composition of control diets may explain the diverse results obtained among studies. Improvement in growth due to seaweed inclusion was also noted in fish by Hashim and Mat Saat (1992), Nakagawa *et al.* (1984), and Nakagawa *et al.* (1987).

Feed intake

A higher feed intake has been observed in shrimp fed diets supplemented with different inclusion levels of *M. pyrifera* or *Sargassum* (Cruz-Suárez *et al.*, 2000; Suárez-García, 2006; Rivera *et al.*, 2002; Gutiérrez-Leyva, 2006). Feeding shrimp a diet supplemented with 3.3% *Ulva* meal resulted in similar feeding intake as for diets supplemented with *Ascophyllum* and *Macrocystis* meals (Cruz-Suárez *et al.*, 2008; Table 4). It has been reported that some compounds from seaweed extracts, such as amino acids, digalactosyl-diacylglycerol, 6-sulfoquinovosyldiacylglycerol, phosphatidylethanolamine and phosphatidylcholine can act as attractants in pelleted diets for abalone (Sakata and Ina, 1985; Sakata *et al.*, 1991) and dimethyl-beta-propionthein (DMTP) for fish (Segovia *et al.*, 2007). Green algae, particularly the Ulvales, are a good source of dimethyl sulfonyl propionate (DMSP) (Van Alstyne *et al.*, 2001). DMSP has been shown to act as an

attractant, and to give improved growth performance in shrimp (Meng-Qing *et al.*, 2001); DMSP increased feed efficiency as well as feed intake.

Feed conversion ratio (FCR)

Changes in FCR due to inclusion of seaweed have been reported. FCR in *Penaeus monodon* was 14% lower in a diet with 10% *Gracilaria heteroclada* (Peñaflorida and Golez, 1996). In contrast, the inclusion of *Macrocystis* and *Sargassum* meal sometimes tended to increase and other times to diminish the FCR (Cruz-Suárez *et al.*, 2000; Rivera *et al.*, 2002; Suárez-García, 2006; Gutierrez-Leyva, 2006). The FCR obtained with *Ulva* meal was significantly lower than that obtained with *Macrocystis* and *Ascophyllum* (Cruz-Suárez *et al.*, 2008b). The inclusion of *Hypnea cervicornis* and *Cryptonemia crenulata* meals, 39% and 26%, exhibited better feed conversion (1.79:1 and 1.82:1) than diets with lower content (2.04:1 and 2.08:1) ($P < 0.05$) (Da Silva and Barbosa, 2008).

Protein efficiency ratio (PER)

Positive results on protein efficiency ratio due to the inclusion of seaweeds have been found in white shrimp *L. vannamei* (Cruz-Suárez *et al.*, 2000a, see Table 4). The improvement in protein utilization may vary with seaweed species. Shrimp fed *Ulva* diet presented higher PER compared with shrimp fed diets containing *Macrocystis* and *Ascophyllum* meals (Cruz-Suárez *et al.*, 2008b). In snakehead fry (Hashim and Mat-Saat, 1992), abalone (Viera *et al.*, 2005) and rohu (Bindu and Sobha, 2004), improved protein efficiency ratios have been observed with the inclusion of pure carragenan, *Ulva* meal, *H. spinella*, *G. cornea*, *Ulva fasciata*, *Spyridia insignis* and *Sargassum wightii*. In contrast, Nakagawa *et al.* (1997) and Valente *et al.* (2006) concluded that the supplementation of *Gracilaria bursa-pastoris*, *Ulva rigida*, *Gracilaria cornea* and *Ascophyllum* in feeds for sea bass did not modify PER. It has been postulated that algae can increase absorption and assimilation of dietary protein (Yone *et al.*, 1986a,b), or modulate lipid metabolism (Nakagawa *et al.*, 1984, 1987, 1997.)

Table 4. Effect of macroalgae meal inclusion on shrimp performance.

Shrimp species	Algae	Inclusion level %	SGR (% day)	weight gain %	Consum g	FCR	PER	Survival %	Ref.	
<i>P. monodon</i> (0.02, 0.05g) 56 days	<i>K. alvarezii</i>	control	4.98			3.8		87	1	
		3,5,7,10	(5%) 5.21			(5%)3.3		(3%) 93		
	<i>G. heteroclada</i>	control	4.17			3.3		83		
		3,5,7,10	(10%) 5.12			(10%) 3.4		83		
			control	4.17			3.3			83
			5,10	3.7			(10%) 2.9			(10%) 83
		5, 10	3.7			(10%) 2.9		(5%) 87		
<i>P. monodon</i> 60 days	<i>Gracilaria spp.</i>	5 to 30	(0-15%)7.9-8			(0 to 15%)3.1-		(0 to 15%)	2	
		control	(30%) 7.3			15%)3.1-3.5		48-56		
<i>L. vannamei</i> (0.6 g) 28days	<i>M. pyrifera</i> Chile	control	2.94	128	2.12a	2.63	1.11	100		
		4	3.12	139	2.49b	2.8	1.04	100		
		8	3.27	151	2.99c	3.12	0.96	100		
(.45g) 28 days	<i>M. pyrifera</i> Mexico	control	4.74a	279.2a	1.68a	1.36a	2.70b	97	3	
		2	5.93b	427.3b	4.07b	2.03b	1.79a	100		
		4	6.18b	470.0b	4.32b	2.04b	1.74a	95		
(.53 g) 28 days	<i>M. pyrifera</i> Mexico	control	3.4	157.6a	1.40a	1.68	1.37	98		
		3.2	3.3	150.6a	1.47a	1.85	1.83	100		
		3.2	3.9	195.0b	2.00b	1.94	1.98	98		
<i>L. vannamei</i> (400mg)	<i>M. pyrifera</i>	control		76a	5	1.9b		100		
		10		100b	6.6	1.7a		100		
		15		75a	6.9	2.3b		86		
		20		62a	6.6	2.8b		77		
<i>L. vannamei</i> (0.16g) 28 days	<i>Sargassum sp.</i>	0		309a	.83a	1.8a		62bc	4	
		2		468bc	1.26 b	1.9a		50ab		
	<i>M. pyrifera</i>	4		561c	1.26 b	1.8a		45a		
		4		404ab	1.29 b	2.2a		52ab		
		control		402ab	1.12 b	2.0a		72c		
<i>F. californiensis</i> (0.5 g) 45 days	<i>Sargassum spp.</i>	0 control	3.64		1.03	1.78		70	5	
		4	3.49		1.15	1.7		77		
<i>L. vannamei</i> (0.34g) 30days	<i>G. cervicornis</i> (Rhodophyta)	0	5.11	363.81		1.56		100	6	
		100	0.44	16.06		30		40		
		50	4.71	313.93		1.84		100		
<i>L. vannamei</i> (1.6 g) 28 days	<i>U. clathrata</i>	3.3		203	5.5	1.71b	1.99b	95	7	
		<i>A. nodosum</i>			187	5.7	1.92ab	1.70a		100
			<i>M. pyrifera</i>			169	5.6	2.1a		1.54a
<i>L. vannamei</i> (1.1 g) 45days	<i>M. pyrifera</i>	control	3.5		1.8ab	2.1a	1.5ab	93.3a	8	
		1	3.8		2.05ab	1.9ab	1.6ab	90a		
		4	3.6		1.81ab	1.9ab	1.6ab	100a		
		7	3.6		1.58 b	1.6b	1.9a	96.7a		
		10	3.9		2.17 ab	1.8ab	1.7ab	86.7a		
	<i>Sargassum sp</i>	1	3.5		1.79 ab	1.9ab	1.6ab	96.7a		
		4	3.9		1.93 ab	1.7ab	1.8ab	93.3a		
		7	3.7		1.85 ab	1.8ab	1.7ab	86.7a		
		10	3.9		2.57 a	2.1a	1.4b	96.7a		
<i>L. vannamei</i> 10-day-old post-larvae 45 days aged	<i>Hypnea cervicornis and Cryptonemia crenulata</i>	39	5.65a			1.8b		96.2a	9	
		26	5.68a			1.8b		97a		
		13	5.17a			2.0a		97a		
		0	4.68a			2.1a		95.2a		

1) Peñaflorida *et al.*, 1996; 2) Briggs and Funge, 1996; 3) Cruz-Suárez *et al.*, 2000; 4) Suárez-García, 2006; 5) Casas-Valdez *et al.*, 2006; 6) Marinho-Soriano *et al.*, 2007; 7) Cruz- Suárez *et al.*, 2008b; 8) Gutiérrez-Leyva, 2006; 9) Da Silva and Barbosa, 2008.

Digestibility

The protein and dry matter digestibility of diets supplemented with low kelp meal concentration, was enhanced in one study and in others it was diminished (Table 5).

Table 5. Apparent digestibility of diets supplemented with seaweed meals

Shrimp species	Algae	Inclusion	APD	ADMD	Reference
		level	%	%	
<i>L. vannamei</i> 3.8 g	<i>M. pyrifera</i> Mexico	control	80.3a	67.9a	Cruz-Suárez <i>et al.</i> , 2000
		2	82.4b	74.1b	
		4	82.9b	77.5c	
<i>L. vannamei</i> 9-14 g	<i>Sargassum sp.</i>	0	88a	78.2a	Suárez-Garcia, 2006
		2	84.9a	71.4a	
		4	86.7a	77.8a	
		4	84.5a	75.1a	
		control	88.1a	81.3a	
<i>L. vannamei</i> 5-7g	<i>Sargassum sp.</i>	0	81.4a	69.6ab	Gutiérrez -Leyva, R.2006
		4	79.7ab	68.9ab	
		10	80.5a	69.1ab	
		4	81.1a	70.1a	
		10	73.3b	60.0b	

Survival and disease resistance

Most of the studies with macroalgae meal supplementation reported excellent survival in shrimp at inclusion levels lower than 10% of the diet.

Peñaflorida and Golez (1996) observed better survival in *P. monodon* fed low inclusion levels of *K. alvarezii* and *G. heteroclada* and increased mortality with increasing levels of these algae. Cruz-Suárez *et al.* (2000) and Suárez-Garcia (2006) also reported excellent survival in shrimp *L. vannamei* fed diets containing (2-4%) kelp meal *M. pyrifera* and *Sargassum spp.* Da Silva and

Barbosa (2008) also reported high survival with diets containing high levels (39, 26 and 13 %) of *Hypnea cervicornis* and *Cryptonemia crenulata* meals. Hashim and Mat-Saat (1992) found a significant improvement in survival of snakehead fry fed a diet supplemented with pure carragenan, followed by diets supplemented with *Polycavernosa spp* meal, *Gracilaria* meal and *Ulva* meal, while the inclusion of *Sargassum* meal resulted in a lower survival.

Several studies reported that dietary supplementation with seaweed meal or their extracts, due to the presence of some compounds (such as fucoidan, alginates, laminarins, carrageenans, etc.) can enhance the immune resistance and improve survival when shrimp are challenged with some bacteria or virus (Table 6).

Table 6. Studies on the immunostimulatory, antibacterial or antiviral effect of seaweed meal or their extracts on penaeid shrimp.

Producto	shrimp species	administration	Seaweed species	Results
Fucoidan	<i>P. japonicus</i> ¹	Oral	60 and 100 mg fucoidan from <i>C. okamuranos</i> semipure/ kg/day	Control WSSV, 77% de survival
Seaweed meal	<i>P. vannamei</i> ^{2,13}	Oral	<i>M. pyrifera</i> meal 1-4% in diets	Partial control WSSV
Fucoidan	<i>P. monodon</i> ⁴	Oral	<i>Sargassum polycystum</i>	Reduce the impact of WSSV
Alginates	<i>L. vannamei</i> ⁵	injected intramuscularly	Sodium Alginate (10, 20 o 50 µg/g)	Increased immune ability as well as resistance to <i>V. alginolyticus</i> infection. Phenoloxidase activity fagocytic activity, bacterial clearance increased significantly when shrimp were injected.
Ulva secondary metabolites	<i>P. monodon</i> ⁶	Oral	<i>Ulva fasciata</i> diet	Increase defense factors such haemogram, agglutination index, phagocytic rate, bacterial clearance and serum bactericidal activity
Hot water extract	<i>L. vannamei</i> ⁸	injected intramuscularly	<i>Gracilaria tenuistipitata</i> (4 o 6 µg/g)	Increased immune ability as well as resistance to <i>V. alginolyticus</i> infection. Total hemocyte count, phenoloxidase activity and respiratory burst increased significantly when shrimp were immersed in seawater containing the extracts or when shrimp were injected.
Seaweed extracts	<i>P. indicus</i> ⁹	injected intramuscularly	<i>C. racemosa</i> , <i>D. dichotoma</i> , <i>E. compressa</i> , <i>G. crassa</i> , <i>G. edulis</i> , <i>H. clathraus</i> , <i>H. musciformis</i> , <i>P. boergeseni</i> , <i>S. wiegthi</i> y <i>T. conoides</i>	Some extracts inactivated partialy WSSV
Hot water extract	<i>P. vannamei</i> ¹⁰	Immersion injected intramuscularly	<i>Sargassum duplicatum</i> (100, 300 y 500 mg/L) immersed or inyected with hot water extract (2, 6, 10 y 20 µg/g).	Increased immune ability as well as resistance to <i>V. alginolyticus</i> infection. Total hemocyte count, phenoloxidase activity and respiratory burst increased significantly when shrimp were immersed in seawater containing the extracts or when shrimp were injected.
Fucoidan extracted three times with 0.05 N HCl at 95 °C for 12 h.	<i>P. monodon</i> ¹¹	injected intramuscularly	<i>Sargassum polycystum</i>	Produced a significantly increased expression (p<0.05) of the ribosomal protein L26 (RPL26) gene, a macrophage activator gene in <i>P. monodon</i>
Fucoidan	<i>P.</i>	oral	<i>Cladosiphon</i>	Partial control against WSSV

Producto	shrimp species	administration	Seaweed species	Results
	<i>vannamei</i> ^{1,2}		<i>okamurnus</i>	infection, variable response

¹Takahashi *et al.*, 1998; ²Cruz-Suárez *et al.*, 2002a; ³Hennequart *et al.*, 2004; ⁴Chotigeat *et al.*, 2004; ⁵Cheng *et al.*, 2004; ⁶Selvin *et al.*, 2004; ⁷Immanuel *et al.*, 2004; ⁸Hou and Chen, 2005; ⁹Balasubramanian *et al.*, 2006; ¹⁰Yeh *et al.*, 2006; ¹¹Deachamag *et al.*, 2006; ¹²Cruz-Suárez *et al.*, 2007a.

Pigmentation

Crustacean pigmentation is affected by dietary pigment source, dosage level, duration of feeding, dietary composition, degree of carotenoid esterification, etc. (Meyers and Latscha, 1997). Menasveta *et al.* (1993) evaluated the pigmentation efficiency of feeds supplemented with 50 ppm astaxanthin vs 5% of extracted brown algae (*Chnoospora minima*) on *P. monodon*, and they found that carotenoides present in the brown algae feed increased carotenoid content in prawn, but the deposition levels were 2 to 3 times lower than those produced by the axtaxanthin supplemented feed. Cruz- Suárez *et al.* (2008b) showed that shrimp fed diets supplemented with 3.3% *Ulva clathrata* meal were highly pigmented in comparison with shrimps fed *Macrocystis* and *Ascophyllum* meals. The pigments present in *Ulva clathrata* (80% lutein) seem to be better metabolized and deposited in shrimp carcass than oxidized forms such as fucoxanthin present in kelp meals, which is in line with the observations reported by Meyers and Latcha (1997) on crustacean carotenoid metabolism. *Ulva* pigmentation efficiency is further better when shrimp are fed with the live co-cultured algae (Cruz-Suárez *et al.*, 2008a).

Brown algae are rich in carotenoids especially in fucoxanthin, and some chlorophyll α and χ , β -carotene and xanthophylls violaxanthin (Barret and Anderson, 1980; Strand *et al.*, 1998; Burtin, 2003; Dhargalkar and Kavlekar, 2004), while green algae, such as *Ulva sp.*, *Ulva clathrata* and *Chaetoniomorpha torta*, contain chlorophyll α and β , β -carotene, lutein, violaxanthin, antheraxanthin, zeaxanthin and neoxanthin (Dhargalkar and Kavlekar, 2004; Burtin, 2003). Red algae contain red phycoerythrin and blue phycocyanin, β -carotene and α -carotene and their dihydroxylated derivatives: zeaxanthin and lutein (Burtin 2003).

Carcass composition

Cholesterol and lipid carcass composition of shrimp can be reduced with the consumption of *Sargassum* meal (Casas-Valdéz *et al.*, 2006) and *Ulva clathrata* (Cruz-Suárez, 2008a). *Ulva* is known to change fat deposition and mobilization patterns in sea bream (Nakagawa *et al.*, 1987), apparently resulting in a more efficient use of fat deposits so weight loss during winter stress is reduced, as well as other compositional changes. The effect may be due at least in part to cysteinolic acid, a non-protein amino acid similar to taurine. Cysteinolic acid, like taurine, can form conjugates with cholesterol in the formation of bile salts, at least in sea bream (Une *et al.*, 1991), and this may be the basis for its effects on fat and cholesterol metabolism. Conceivably, animals naturally adapted to diets that include cysteinolic acid require it for efficient fat metabolism. Cysteinolic acid could play other roles in metabolism as well. For example, it could be converted to cysteine.

Single cell seaweed detritus as hatchery diets for shrimp

The use of seaweeds detritus as larvae feed has been reported by Japanese researchers. The formation of algal detrital particles in a completely cell-detached form, namely, single cell detritus (SCD) was reported for the first time by Uchida *et al.* (1999) during the microbial degradation process of macroalgal thalli. SCD displays 3 characteristics suitable for feed for aquatic hatchery animals: 1) The size of SCD is in the 2-14 µm range, which is similar to that of dietary phytoplankton; 2) The cell wall components of SCD are partially degraded which facilitates digestion; 3) Bacterial cells are attached to SCD, which modifies the algal detritus to protein-rich particles. The use of SCD as a potential hatchery diet instead of phytoplankton culture, which is labor-intensive, was successfully demonstrated based on feeding experiments with *Artemia*. Further modification of SCD could include the attachment of bacteria to SCD which would exert beneficial effects on the hatchery animals. Use of SCD diets in fish feeding regimes is an attempt to introduce the concept of detrital food web to aquaculture systems, which could contribute to the development of sustainable fish nursery systems (Uchida *et al.*, 1997; Uchida and Murata, 2002).

A technology to produce shrimp hatchery diets from brown and green macroalgae was developed and patented by Uchida and Numaguchi (1996, USA patent 5801050). “Algal detritus particles are prepared by contacting algae with marine bacteria capable of attaching to and decomposing the algae under conditions sufficient to induce decomposition of the structural components of the algal tissue, thereby forming detritus suitable for use as a primary feed for marine organisms. Marine bacteria which belong to the genus *Alteromonas* can be used. Algae belonging to the class Phaeophyceae or Chlorophyceae preferably are used as the starting material in the invention. *Laminaria*, *Eisenia*, *Ecklonica*, *Undaria* and *Sargassum* are examples of types of Phaeophyceae which are useful. These algae can be decomposed by the above described marine bacteria to form particles suitable for use as a primary feed. *Ulva pertusa*, *Monostroma*, *Enteromorpha* and *Acetabularia* are examples of types of Chlorophyceae. The cell walls of these plants can be decomposed by the action of the above marine bacteria, resulting in formation of protoplasmic detritus suitable for use as primary feed comprising particles having substantially the same diameter. Bacteria which are most preferred for accomplishing the present invention include *Pseudoalteromonas espejiana*. The proliferation of bacteria is faster and requires less use of resources such as light or heat compared with that of phytoplankton or zooplankton, which are used for feed at present. Thus, using the present method, detritus feed can be produced at low cost and with less labor. The invention provides an avenue for utilizing undeveloped or underdeveloped marine resources, such as *Ulva pertusa*, into value-added products in the marine environment. The concept of making use of and/or recycling marine resources has the advantage of preserving the environment” (Uchida and Numaguchi (1996, USA patent 5801050).)

The Fisheries College and Research Institute (FCRI) in India, has also developed a single cell diet that can be used as a cost-effective substitute for micro algae cultures for shrimp larvae reared at marine hatcheries. Named ‘Marine Single Cell Detritus,’ the product was derived from seaweeds. The feed was developed through enzymatic and fermentative treatment of seaweeds in two phases. In the first stage, the seaweeds were treated with an enzyme which led to the formation of single cell units. The ‘enzymatic digest’ was then treated with bacteria and yeast in the ‘fermentative phase’ to form the product. A series of tests conducted by the institute proved that MSCD had distinct advantages over traditional feeds like micro algae cultures and imported feeds like ‘artemia cysts,’ used widely in marine hatcheries across the country now. The MSCD

is 20% less expensive than artemia cysts and its manufacturing process is simpler than that of micro algae cultures. Besides, the new diet possesses bioremediation properties which control water quality. The 'probiotic' characteristics of the feed helps the fish develop infection resistance. The MSCD has 35% crude protein, making it nutritious. It can be stored in room temperature for a year

([http://www.hinduonnet.com/thehindu/thscrip/p...007073157820100.htm&date=2007/07/31/&prd=th&\(1 of 2\)](http://www.hinduonnet.com/thehindu/thscrip/p...007073157820100.htm&date=2007/07/31/&prd=th&(1%20of%202))).

Fresh algae and shrimp/macroalgae co-culture

There are few reports about the nutritional benefits that shrimp can get as a result of fresh algae consumption or about shrimp/macroalgae integrated cultures (Table 7).

The use of *Enteromorpha sp* as part of fresh food to induce maturation of prawn *Penaeus indicus* and *L. stylirostris* was reported by Emerson (1980), Emerson *et al.* (1983), and Bray and Lawrence, (1988, 1990) respectively. Moss (1994) studied the nutritional contribution of different macro- and microalgae to juvenile shrimp growth. Shrimp juvenile white shrimp, *Penaeus vannamei* fed a diatom culture composed primarily of *Chaetoceros sp.* were significantly heavier ($p < 0.05$) than shrimp fed a monoculture of the green alga *Nannochloropsis oculuta*, fronds from the leafy macroalga *Ulva sp.*, or fronds from the filamentous macroalga *Enteromorpha sp.* after 5 days. Porchas-Cornejo *et al.* (1999) reported that shrimp *Farfantepenaeus californiensis* increases its growth rate 3 fold in the presence of the algae *Caulerpa sertularioides*. Lombardi *et al.* (2006) tested the feasibility of co-culturing the Pacific white shrimp *Litopenaeus vannamei* and the Philippines seaweed *Kappaphycus alvarezii* in floating cages; juveniles (2.39 ± 1.62 g) were transferred to 6 experimental grow-out cages at a density of 100 shrimp/m². Commercial pellets consisting of 35% crude protein were supplied once a day and shrimp were harvested after 103 days of growth. Thalli of the seaweed *K. alvarezii* were fixed on ropes and attached to floating tubes and set up inside 3 shrimp grow-out cages. Shrimp yield reached production rates as high as 3.23 kg/m²/yr with a mean of 2.36 ± 0.76 kg/m²/yr. Seaweed production reached rates of 23.70 kg/m²/yr with a mean of 17.47 ± 5.71 kg/m²/yr. There were no negative interferences in co-culturing shrimp and seaweed inside the

same cage. Student's t-test showed no statistical differences between the two treatments (monoculture and polyculture) for shrimp weight gain, survival rate, and feed conversion ratio.

Van Tri and Thi Thanh Ha (2004) evaluated 3 diet formulas: **live seaweed** + mixed feed; dry seaweed + mixed feed; and dry seaweed on N5 Nauplius *Litopenaeus vannamei* of the same size-collected from a hatchery at a density of 90 individuals/liter. The authors showed that white leg shrimp larvae feeding on live seaweed and mixed feed have shorter metamorphosis period than those feeding on dry seaweed and mixed feed or dry seaweed only. Larvae fed on live seaweed together with mixed feed had higher growth rate (in length and weight) and higher survival rate (48-53%) than those fed on dry seaweed and mixed feed or dry seaweed only.

The use of *Ulva* as food for shrimp in co-culture experiment has been proved (Cruz-Suárez *et al.*, 2008a) to be beneficial as an alternative to improve the utilization of commercial feed and increase the economical value of the shrimp produced due to a larger size and a higher pigmentation. Shrimps in co-culture groups fed *Ulva* and pelleted feed showed significantly better performance than the shrimp fed *Ulva* or feed alone. The consumption of co-cultivated *Ulva clathrata* by shrimp *L. vannamei* improved the pelleted feed utilization and the growth rate: with 10 to 45% less commercial feed, growth rate was improved by 60%. Additionally, shrimp carcass quality improved with the level of *Ulva* consumed: lipids diminished, while total carotenoides, total and esterified astaxanthin increased following quadratic models. The co-cultivated *Ulva clathrata* also modified water quality: decreasing water turbidity and total phytoplankton cell number (Cruz-Suárez *et al.*, 2008a).

In the shrimp/macroalga integrated system, nitrogenous enriched waste water of cultured shrimp may be transformed into a valuable algal biomass, seaweeds production being an added income as feed for shrimp (Evans and Langdon, 2000; Schuenhoff *et al.*, 2003; Neori *et al.*, 2004). Several studies have shown that culture of *Ulva spp.* in nutrient-rich waters increases its protein content from 11% to over 32% in dry weight (Shpigel *et al.*, 1999; Boarder and Shpigel, 2001). This biofilter produced *Ulva* has been shown to provide good growth rates for abalone *H. tuberculata* (Neori *et al.*, 1998; Shpigel *et al.*, 1999), *H. discus hannai* (Corazani and Illanes, 1998; Shpigel *et al.*, 1999) and *H. roei* (Boarder and Shpigel, 2001). The evaluation of the

suitability of macroalgae cultivated in an integrated biofilter unit as a potential feed for the shrimp is a sustainable alternative to diminish shrimp production cost and environmental impact.

Polyculture of seaweeds in shrimp ponds or in the effluent ponds is one of the Best Management Practices suggested by Food and Agriculture (FAO) experts.

Table 7. Studies on the use of fresh algae as feed and shrimp/macroalgae co-culture

Shrimp species	Title	References
<i>P. indicus</i>	Induced maturation of prawn <i>Penaeus indicus</i> . (<i>Enteromorpha sp.</i> Fresh)	Emerson, W. D. 1980.
<i>P. indicus</i>	Growth and Maturation of <i>Penaeus indicus</i> under blue and green light. (<i>Enteromorpha sp.</i> Fresh)	Emerson, W. D., Hayes, D.P., and Ngonyame, M., 1983.
<i>Litopenaeus vannamei</i>	Growth rates, nucleic acid concentrations, and RNA/DNA ratios of juvenile white shrimp, fed different algal diets. <i>Chaetoceros sp.</i> green algae <i>Nannochloropsis oculuta</i> , fronds from the leafy macroalga, <i>Ulva sp.</i> , or fronds from the filamentous macroalga, <i>Enteromorpha sp.</i> after 5 days.	Moss, 1994
<i>P. stylirostris</i>	Reproduction of eyestalk-ablated <i>Penaeus stylirostris</i> fed various levels of total dietary (<i>Enteromorpha sp.</i> Fresh)	Bray <i>et al.</i> , 1990a
<i>P. monodon</i>	Influence of dietary fatty acids on reproduction of <i>P. stylirostris</i> (<i>Enteromorpha sp.</i> Fresh)	Bray and Lawrence, 1988.
<i>Penaeus californiensis</i>	Efecto de la macrolaga <i>Caulerpa sertularioides</i> en el desarrollo del camaron café	Porchas Cornejo <i>et al.</i> , 1999
<i>Litopenaeus vannamei</i>	Impacts of feed on the growth of white leg shrimp larvae <i>Litopenaeus vannamei</i> (Boone,1931).	Van and Thanh , 2004.
<i>Litopenaeus vannamei</i>	Cage polyculture of the Pacific white shrimp and the Philippines seaweed <i>Kappaphycus alvarezii</i>	Lombardi <i>et al.</i> , 2006.
<i>Litopenaeus vannamei</i>	Shrimp and green algae <i>Ulva clathrata</i> co-culture to optimize commercial feed utilization.	Cruz- Suárez <i>et al.</i> , 2008a.
<i>Litopenaeus vannamei</i>	Cultivation of seaweed <i>Ulva clathrata</i> within shrimp ponds	Tomena and Copertino XIX ISS 2007 (214)
<i>Litopenaeus vannamei</i>	Biofiltering efficiency, uptake and assimilation rates of <i>Ulva clathrata</i> cultivated in shrimp aquaculture waste water	Copertino and Tomena XIX ISS 2007 (125)
<i>P. monodon</i>	Possible implications of the co-cultivation of blacktiger shrimp and cladophoraceae species on southwest asin shrimp farm	Tsutsui <i>et al.</i> , XIX ISS 2007 (66)
	Development of abandoned shrimp ponds for restoration of wild <i>Gracilaria</i> along the coast of Pattan Bay, south of Thailand	Rapepom <i>et al.</i> , XIX ISS 2007 (37)
<i>P. monodon</i>	Profitable environmental remediation with seaweeds in an intensive marin shrimp culture	Hamano <i>et al.</i> , . XIX ISS 2007 (232)

Conclusions

Algal meal inclusion in aquafeeds generally improves pellet stability and pellet water absorption. Growth performance, FCR, PER and disease resistance generally are improved at low inclusion levels. Additionally, shrimp product quality may also be improved by a better body pigmentation and lower lipid and cholesterol content. The use of fresh seaweeds for brood stock and fermented seaweeds for larvae feeds seems to be an interesting and promising alternative for these shrimp life stages. The shrimp/algae co-culture using live algae as a suitable feed for shrimp represent a sustainable alternative to diminish the need for artificial feed. The benefit of co-culture is amplified by the fact that co-cultured algae are enriched in highly available nutrients and variations in composition may be less than in the wild.

References

- Ackman, 1981 Alga as sources of edible oils in new sources of fat and oils. Pryde E.H., Princen L.H., and Mujerhee K.D. Ed. *New Sources of Fats and Oils: [contributions presented at a symposium at the International Society for Fat Research/American Oil Chemists' Society World Congress, April 27-May 1 1980 in New York City]* International Society for Fat Research, American Oil Chemists' Society. The American Oil Chemists Society, 340 pp (pp189 a 220)
- Aguilera-Morales, M., Casas-Valdez, M., Carrillo-Dominguez, S., Gonzalez-Acosta, B., Pérez-Gil, F. (2005) Chemical composition and microbiological assays of marine algae *Enteromorpha spp.* as a potential food source. *Journal of Food Composition and Analysis*, 79-88.
- Balasubramanian, G., Sudhakaran, R., Syed-Musthaq, S., Sarathi, M. & Sahul-Hameed, A.S. (2006) Studies on the inactivation of white spot syndrome virus of shrimp by physical and chemical treatments, and seaweed extracts tested in marine and freshwater animal models. *J Fish Dis.*, 29(9), 569-572.
- Barrett, J. & Anderson, J.M. (1980) The P-700-chlorophyll alpha-protein complex and two major light-harvesting complexes of *Acrocarpia paniculata* and other brown seaweeds. *Biochim Biophys Acta.*, 590(3), 309-323.
- Bindu, M.S. & Sobha, V. (2004) Conversion efficiency and nutrient digestibility of certain seaweed diets by laboratory reared *Labeo rohita* (Hamilton). *Indian J Exp Biol.*, 42(12), 1239-1244.
- Bjerkeng, B. (2000) Carotenoid pigmentation of salmonid fishes- Recent progress. pp. 71-89 In *Avances en nutrición Acuicola V. Memorias del V symposium Internacional de Nutrición Acuicola* (Cruz-Suárez L. E., Rique-Marie D., Tapia-Salazar M., Olvera-Novoa M. A. & Civera-Cerecedo R. Eds.). Universidad Autónoma de Nuevo Leon.
- Boarder, S.J. & Shpigel, M. (2001) Comparative performances of juvenile *Haliotis roei* fed on enriched *Ulva rigida* and various artificial diets. *J. Shellfish Res.*, 20 (2), 653-657.
- Boonyaratpalin, M., Thongrod, S., Supamattaya, K., Britton, G. & Schlipalius, L.E. (2001) Effects of β -carotene source, *Dunaliella salina*, and astaxanthin on pigmentation, growth, survival and health of *Penaeus monodon*. *Aquaculture Research*, 32(1), 182-190.
- Bray W.A. and Lawrence A.L., 1988. Influence of dietary fatty acids on reproduction of *P. stylirostris* (abstract) J. World. Aquac. Soc. 19:18A
- Bray W. A., Lawrence, A.L. and Lester, L.J., 1990a. Reproduction of eyestalk-ablated *Penaeus stylirostris* fed various levels of total dietary. J. World Aquacul. Soc. 21: 41-52
- Briggs, M.R.P. and Funge-Smith, S-l., 1996. The potential of *Gracilaria spp.* meal for supplementation of diets for juvenile *Penaeus monodon* Fabricius. *Aquaculture Research* 27, 345-354.
- Burtin, P. (2003) Nutritional value of seaweeds. *Electronic Journal of Environmental, Agricultural and Food Chemistry*, 2(4), 498-503.
- Castro-González, M.I., Carrillo, S., Pérez-Gil, F., Manzano, R. & Rosales, E. (1991) *Macrocystis pyrifera*: Potential resource for animal feeding. *Cuban Journal of Agricultural Science*, 25(1), 77-81.

- Castro-González, M.I., Carrillo-Domínguez, S. & Pérez-Gil, F. (1994) Chemical composition of *Macrocystis pyrifera* (Giant Sargazo) collected in summer and winter and its possible use in animal feeding. *Ciencias Marinas*, 20(1), 33-40.
- Casas-Valdez M, Hernández C., Aguila R., González B., Marín A., Rodríguez S., Carrillo S., Pérez Gil F., Cruz-Suárez L.E., Ricque D. & Tapia M. 2002. *Sargassum spp.* como fuente potencial de alimento para camarón. Informe Técnico Final. CGPI. Instituto Politécnico Nacional, 34 pp.
- Casas-Valdez, M., Portillo-Clark, G., Aguila-Ramírez, N., Rodríguez-Astudillo S., Sánchez-Rodríguez I. y Carrillo-Domínguez S. 2006 Efecto del alga marina *Sargassum spp.* sobre las variables productivas y la concentración de colesterol en el camarón café, *Farfantepenaeus californiensis* (Holmes, 1900). *Rev. biol. mar. oceanogr.* [online]. jul., vol.41, no.1 , p.97-105.
- Cerecer-Cota, E.E., Ricque-Marie, D., Mendoza-Cano, F., Nieto-Lopez, M. & Cruz-Suárez, L.E. (2005) Pellet stability and hardness influence feed consumption of Pacific White Shrimp. *Global Aquaculture Advocate*, (2), 85-86.
- Cheng, W., Liu, C.H., Yeh, S.T. & Chen, J.C. (2004) The immune stimulatory effect of sodium alginate on the white shrimp *Litopenaeus vannamei* and its resistance against *Vibrio alginolyticus*. *Fish Shellfish Immunol.* 17(1), 41-51.
- Chotigeat, W., Tongsupa S., Supamataya, K. & Phongdara, A. (2004) Effect of Fucoïdan on Disease Resistance of Black Tiger Shrimp. *Aquaculture*, 233(1-4), 23-30.
- Corazani, D., Illanes, J.E., 1998. Growth of juvenile abalone, *Haliotis discus hannai* Ino 1953 and *Haliotis rufescens* Swainson 1822, fed with different diets. *J. Shellfish Res.*, 17(3), 663– 666.
- Cruz-Suárez, L.E., Ricque-Marie, D., Tapia-Salazar, M. & Guajardo-Barbosa, C. (2000) Uso de harina de kelp (*Macrocystis pyrifera*) en alimentos para camarón. p. 227-266. Editores L.E. Cruz-Suárez, Denis Ricque-Marie, Mireya Tapia-Salazar, Miguel A. Olvera-Novoa, y Roberto Cerecedo-Olvera. Avances en Nutrición Acuicola V. Memorias del Quinto Simposium Internacional de Nutrición Acuicola. 19-22 Noviembre. Universidad Autónoma de Nuevo León. Monterrey, Nuevo León, México. ISBN 970-694-52-9.
- Cruz-Suárez, L.E., Ricque-Marie, D., Tapia-Salazar, M., Guajardo Barbosa, C., Cleveland, K. & Hasson, K. (2002a) Efecto de alimentos con harina de kelp en la prevención de infección viral (WSSV) en camarón blanco *L. vannamei*. *Panorama Acuicola*, 7(4), 16-19.
- Cruz-Suárez, L.E., Ricque Marie, D., Tapia Salazar, M., Guajardo Barbosa, C., Obaldo, Leonard, Velasco-Escudero, M. & Carrasco, A. (2002b) Water stability and texture of shrimp pelleted feeds formulated with natural and synthetic binders. *Global Aquaculture Advocate* 6, 44-45.
- Cruz- Suárez et al., 2003. Uso de alga *Sargassum sp.* en la elaboración de dietas para camarón y su impacto en los parámetros productivos XIII Congreso Latinoamericano de Nutrición. Acapulco, México. 9-13 de Noviembre
- Cruz-Suárez L.E., Ruíz-Díaz, P.P., Cerecer Cota E.E., Nieto-Lopez, M., Guajardo-Barbosa C., Tapia-Salazar, M., Ricque-Marie D. (2006a). Revisión sobre características físicas y control de calidad de alimentos comerciales para camarón en México. 330-370 pp . Eds: *L. Elizabeth Cruz- Suárez, Denis Ricque-Marie,*

Mireya Tapia Salazar, Martha G. Nieto López, David A. Villarreal Cavazos, Ana C. Puello Cruz y Armando García Ortega. Avances en nutrición acuícola VIII. Memorias del VIII Simposio Internacional de Nutrición Acuícola. Universidad Autónoma de Nuevo León. Monterrey, N.L. México. ISBN 970-694-331-5.

- Cruz-Suárez, L. E., Hernández, J., Porchas-Conrnejo, M., Coronado-Molina, D.E., Linné-Unzueta-Bustamante, M.L., Nieto-López, M., Tapia-Salazar, M. & Ricque-Marie, D. (2006b). Estudio del efecto antiviral del fucoïdan del alga café *Cladosiphon okamuranus* contra la infección por WSSV en camarón blanco *Litopenaeus vannamei*. p. 78. L. Elizaebth Cruz Suárez, Denis Ricque-Marie, Mireya Tapia Salazar, Martha G. Nieto López, David A. Villarreal Cavazos, Ana C. Puello Cruz y Armando García Ortega. Book proceedings of the VIII simposio internacional de Nutricion acuicola. 15 to 17 of November 2006. Universidad Autónoma de Nuevo León. Monterrey, N.L. México.
- Cruz-Suárez, L. E., Hernández, J., Porchas-Cornejo, M., Coronado-Molina, D. E., Linné-Unzueta- Bustamante, M. L., Nieto-López, M., Tapia-Salazar, M.L., Ricque-Marie, D. 2007a. The antiviral effect of algal fucoïdians. Fucoïdan, extracted from the brown algae *Cladosiphon okamuranus* has been shown to be effective against WSSV in the white shrimp *Litopenaeus vannamei*. AQUA Culture AsiaPacific Magazine January/February 2007
- Cruz-Suárez, L.E. and Tapia-Salazar M., 2007b. Harina de Kelp. En Manual de Ingredientes proteicos y aditivos empleados en la formulación de alimentos balanceados para camarones peneidos. Ed. Tsai Garcia Galano, Humberto Villarreal Colmenares y Jorge L. Fenucci.
- Cruz-Suárez L.E., León A.A., Peña-Rodríguez A., Rodríguez-Peña G., Moll B. , Ricque-Marie D. 2008a. Shrimp and green algae co-culture to optimize commercial feed utilization. ISNF XIII International Symposium on Nutrition and Feeding in Fish. Florianopolis. June 1 to 5, Brazil
- Cruz-Suárez, L.E., M. Tapia-Salazar, M.G. Nieto-López, C. Guajardo-Barbosa & D. Ricque-Marie 2008b., Comparison of *Ulva clathrata* and the kelps *Macrocystis pyrifera* and *Ascophyllum nodosum* as ingredients in shrimp feeds. Aquaculture Nutrition. Volume 9999, Issue 9999 , Pages - 2007 Blackwell Publishing Ltd.
- Da Silva R.L. da & Barbosa J. M. (2008). Seaweed meal as a protein source for the white shrimp *Litopenaeus vannamei*. J Appl Phycol
- Deachamag, P., Intaraphad, U., Phongdara, A. & Chotigeat, W. (2006) Expression of a Phagocytosis Activating Protein (PAP) gene in immunized black tiger shrimp. *Aquaculture*, 255 (1-4), 165-172.
- Dhargalkar, V.K. & Kavlekar, D. (2004) Seaweeds- a field manual. Verlecar X. N. & Devanand Kavlekar (Eds.). National Institute of Oceanography, Dona Paula, Goa. 403 004. 1st edition. 42 pp. <http://drs.nio.org/drs/bitstream/2264/96/1/Seaweeds-Manual.pdf>.
- Emerson, W. D. 1980. Induced maturation of prawn *Penaeus indicus*. Mar. Ecol. Prog. Ser. 2:121-131
- Emerson, W. D., Hayes, D.P., and Ngonyame, M., 1983. Growth and Maturation of *Penaeus indicus* under blue and green light. S Africa. J. Zool. 18: 71-75
- Evans, F., Langdon, C.J., 2000. Co-culture of dulse *Palmaria mollis* and red abalone *Haliotis rufescens* under limited flow conditions. *Aquaculture*, 185, 137– 158.

- Fleurence, J. (1999). Seaweed proteins: biochemical nutritional aspects and potential uses. *Trends in Food Science & Technology*, 10(1), 25–28.
- Gutiérrez-Leyva, R. 2006. Uso de harinas de *Macrocystis pyrifera* y *Sargassum* spp. en alimentos balanceados para el camarón *Litopenaeus vannamei*: efectos sobre el crecimiento y la digestibilidad *in vivo*. Tesis maestría en ciencias. CICIMAR-IPN La Paz, Baja California Sur. México. 84 p.
- Hashim, R. & Mat-Saat, N. A. (1992) The utilization of seaweed meals as binding agents in pellet feeds for snakehead (*Channa striatus*) fry and their effects on growth. *Aquaculture*, 108, 299-308.
- Hennequart, F., O'Connell, E.O., Tuohy, J.S. & Tuohy, M.G. (2004) Brown macro-algae. *Aquafeeds: Formulation and Beyond*, 1(4), 14-18.
- Hou, W.Y. & Chen, J.C. (2005) The immunostimulatory effect of hot-water extract of *Gracilaria tenuistipitata* on the white shrimp *Litopenaeus vannamei* and its resistance against *Vibrio alginolyticus*. *Fish Shellfish Immunol.*, 19(2), 127-138.
- Immanuel, G., Vincybai, V.C., Sivaram, V., Palavesam, A., Marian, M.P., 2004. Effect of butanolic extracts from terrestrial herbs and seaweeds on the survival, growth and pathogen (*Vibrio parahaemolyticus*) load on shrimp *Penaeus indicus* juveniles. *Aquaculture* 236, 53– 65
- Jimenez-Escrig, A. & Sanchez-Muniz, F.J. (2000) Dietary fibre from edible seaweeds: chemical structure, physicochemical properties and effects on cholesterol metabolism. *Nutrition Research*, 20(4), 585-598.
- Kuda, T., Yokoyama, M. & Fujii, T. (1997) Effects of marine algal diets hijiki, ao Nori, and Nori on levels of serum lipids and cecal microflora en rats. *Fish Sci.*, 63, 428-432.
- Lombardi, J. V., de Almeida Marques H. L., Toledo Lima Pereira R., Saleé Barreto O. J., and de Paula E.J. 2006. Cage polyculture of the Pacific white shrimp *Litopenaeus vannamei* and the Philippines seaweed *Kappaphycus alvarezii*. *Aquaculture* 258, no. 1-4: 412-415.
- Marinho-Soriano, E., Camara M. R., de Melo Cabral T., and do Amaral Carneiro M. A., (2007). Preliminary evaluation of the seaweed *Gracilaria cervicornis* (Rhodophyta) as a partial substitute for the industrial feeds used in shrimp (*Litopenaeus vannamei*) farming. *Aquaculture Research* 38, no. 2: 182-187.
- McHugh, D.J. (2003) A guide to Szeged industry, FAO Fisheries Technical paper, No. 441. Rome, FAO, p 105.
- Mabeau, S., Fleurence, J., 1993. Seaweed in food products: biochemical and nutritional aspects. *Trends Food Sci. Technol.*, 4, 103–107.
- Marais, M.F. & Joseleau, J.P. (2001) A fucoidan fraction from *Ascophyllum nodosum*. *Carbohydr Res.*, 336(2), 155-159.
- Marsham, S., Scott, G.W., Tobin, M.L. (2007) Comparison of nutritive chemistry of a range of temperate seaweeds. *Food chemistry*, 100, 1331-1336.
- Men-qing L., Qing Ch., and Aksnes A. (2001) Identification of feeding stimulants for shrimp. *Marine Fisheries research*. Vol 22 (4) p. 71-74
- Menasveta, P, Worawattanamateekul, W., Latscha, T., Clark, J.S., 1993. Correction of black tiger prawn (*Penaeus monodon* Fabricius) coloration by astaxanthin. *Aquaculture Engineering* 12, 203-213. Meyers, S.P. & Latscha,

- T. (1997). Carotenoids. In: Crustacean Nutrition (D'Ábramo, L.R.D., Conclin, D.E. & Akiyama, D.M. Eds.). Advances in World Aquaculture Vol. 6. The World Aquaculture Society.
- Moss, S.M. (1994) Growth rates, nucleic acid concentrations, and RNA/DNA ratios of juvenile white shrimp, *Penaeus vannamei* Boone, fed different algal diets. *Journal of Experimental Marine Biology and Ecology*, 182, 193-204.
- Nakagawa, H., Kasahara, S., Sugiyama, T. & Wada, I. (1984) Usefulness of *Ulva* meal as feed supplement in cultured black sea bream. *Suisan Zoshoku*, 32, 20-27.
- Nakagawa, H., Kasahara, S., Sugiyama, T. (1987) Effect of *Ulva* meal supplementation on lipid metabolism of black sea bream *Acanthopagrus schlegeli*. *Aquaculture*, 62, 106-121.
- Nakagawa, H., Umino, T. & Tasaka, Y. (1997) Usefulness of *Ascophyllum* meal as feed additive for red sea bream, *Pagrus major*. *Aquaculture*, 151, 275-281.
- Nakagawa, H., and Montgomery W.L. (2007). Algae. In: Nakagawa H., Sato M., and Gatlin III D.M., Dietary supplements for the health and quality of culture fish. CABI, USA. Pag 133-168.
- Neori, A., Ragg, N.L.C. & Shpigel, M. (1998) The integrated culture of seaweed, abalone, fish and clams in modular intensive landbased systems: II. Performance and nitrogen partitioning within an abalone (*Haliotis tuberculata*) and macroalgae culture system. *Aquac. Eng.*, 17 (4), 215–239.
- Neori, A., Chopin, T., Troell, M., Buschmann, A.H., Kraemer, G.P., Halling, C., Shpigel, M., Yarish, C., (2004). Integrated aquaculture: rationale, evolution and state of the art emphasizing seaweed biofiltration in modern mariculture. *Aquaculture*, 231, 361– 391.
- Obluchinskaia, E.D., Voskoboinikov, G.M. & Galynkin, V.A. (2002) Content of alginic acid and fucoidan in fucus algae of the Barents sea. *Prikl Biokhim Mikrobiol.*, 38(2), 213-216.
- Paradossi, G., Cavalieri, F., Pizzoferrato, L. & Liquori, A.M. (1999). A physico-chemical study on the polysaccharide ulvan from hot water extraction of the macroalga *Ulva*. *Int J Biol Macromol.*, 25(4), 309-315.
- Peñaflorida, V.D. & Golez, N.V. (1996) Use of seaweed meals from *Kappaphycus alvarezii* and *Gracilaria heteroclada* as binders in diets of juvenile shrimp *Penaeus monodon*. *Aquaculture*, 143, 393-401.
- Percival, E. (1968) Glucuronyloxylofucan, a cell-wall component of *Ascophyllum nodosum*. *Carbohydrate Research*, 7(3), 272-283.
- Porchas Cornejo M., Martinez Cordova L., Magallon Barajas F. & Portillo Clark G. (1999) Efecto de la macrolaga *Caulerpa sertularioides* en el desarrollo del camarón café *Penaeus californiensis* (Decapoda: Peneidae). *Revista de Biología Tropical* 47,437-442.
- Ray, B. & Lahaye, M. (1995a) Cell-wall polysaccharides from marine green alga *Ulva rigida* (Ulvales, Chlorophyta). Extraction and chemical composition. *Carbohydrate Research*, 274, 252-261.
- Ray, B. & Lahaye M. (1995b) Cell-wall polysaccharides from marine green alga *Ulva rigida* (Ulvales, Chlorophyta). Chemical structure of ulvan. *Carbohydrate Research*, 274, 313-318.

- Rivera, G., F. Yoong, G. Riofrío, B. Reinoso, F. Hurtado & P. Massuh. (2002). Inclusión de harina de kelp (*Macrocystis pyrifera*) en alimentos balanceados para el camarón. Congreso Iberoamericano Virtual de Acuicultura. CIVA. <http://www.civa2002.org>, 244-252.
- Rodríguez-Montesinos, Y.F. & Hernández-Carmona, G. (1991) Variación estacional y geográfica de la composición química de *Macrocystis pyrifera* en la costa occidental de Baja California. *Ciencias Marinas*, 17, 3, 91-107.
- Sakata, K. & Ina, K. (1985) Digalactosyldiacylglycerols and phosphatidylcholines isolated from a brown alga as effective phagostimulants for a young abalone. *Bull. Jpn. Soc. Sci. Fish.*, 51, 659–665.
- Sakata, K., Kato, K., Iwase, Y., Okada, H., Ina, K. & Machiguchi, Y. (1991) Feeding-stimulant activity of algal glycerolipids for marine herbivorous gastropods. *Journal of Chemical Ecology*, 17(1), 185-193.
- Selvin, J., Huxley, A.J., Lipton, A.P. (2004). Immunomodulatory potential of marine secondary metabolites against bacterial diseases of shrimp. *Aquaculture*, vol. 230 (1-4), 241-249.
- Schuenhoff, A., Shpigel, M., Lupatsch, I., Ashkenazi, A., Msuya, F.E., Neori, A., (2003). A semi-recirculating, integrated system for the culture of fish and seaweed. *Aquaculture*, 221, 167– 181.
- Sharp, G. (1987) *Ascophyllum nodosum* and its harvesting in Eastern Canada. In: Case studies of seven commercial seaweed resources (Doty M.S., Caddy J.F. & Santelices B. Eds.). FAO Fisheries Technical paper-281.
- Shpigel, M., Ragg, N.C., Lupatsch, I. & Neori, A. (1999) Protein content determines the nutritional value of the seaweed *Ulva lactuca* for the abalone *Haliotis tuberculata* and *H. discus hannai*. *J. Shellfish Res.*, 18, 223– 227.
- Strand, A., Herstad, O. & Liaaen-Jensen, S. (1998) Fucoxanthin metabolites in egg yolks of laying hens. *Comp Biochem Physiol A Mol Integr Physiol.*, 119(4), 963-974.
- Suárez-García, H.A. (2006) Efecto de la inclusión de alginato y harina de algas *Sargassum sp* y *Macrocystis pyrifera* sobre la estabilidad en agua, digestibilidad del alimento y sobre el crecimiento del camarón blanco *Litopenaeus vannamei*. Undergrate thesis. Universidad Autónoma de Nuevo León, Mexico.
- Suzuki, T., Ohsugi, Y., Yoshie, Y. & Hirano, T. (1996) Dietary fiber content, water holding capacity and binding capacity of seaweeds. *Fish Sci.*, 62, 454-461.
- Takahashi, Y., Uehra, K., Watanabe, R., Okumura, T., Yamashita, T., Omura, H., Kawano T., Kanemitsu, A., Narasaka, H., Suzuki, N. & Itami, T. (1998) Efficacy of oral administration of fucoidan, a sulphated polysaccharide, in controlling white spot syndrome in kuruma shrimp in Japan. In *Advances in shrimp biotechnology* (Flegel T.W. Ed.). National Center for Genetic Engineering and Biotechnology, Bangkok.
- Une M., Goto T., Kihira K., Kuramoto T., Hagiwara K., Nakajima T., and Hoshita T., (1991). Isolation and identification of bile salts conjugated with cysteinolic acid from bile of the red seabream, *Pagrosomus major*. *Journal of Lipid Research*, Volume 32, p. 1619-1623.
- Uchida, M., Nakata, K., Maeda, M., (1997). Introduction of detrital food webs into aquaculture system by supplying single cell algal detritus produced from *Laminaria japonica* as a hatchery diet for *Artemia nauplii*. *Aquaculture*, vol. 154 (2)pp. 125-137
- Uchida M., Nugamuchi, K.,(1998). Method for preparing algal detritus. \united States Patent 5801050

- Uchida, M. and Murata M., (2002). Fermentative preparation of single detritus from seaweed, *Undaria pinnatifida*, suitable as replacement hatchery diet for unicellular algae. *Aquaculture*. Vol. 207 (3-4) pp. 345-357
- Vairappan Ch., and Shigeharu, S., 2006. Probiotic fortified tropical seaweed silage as aquaculture feed. WAS 2006. AQUA 2006/ Meeting Abstract.
- Valente, L.M.P., Gouveia, A., Rema, P., Matos, J., Gomes E.F. & Pinto, I.S. (2006) Evaluation of three seaweeds *Gracilaria bursa-pastoris*, *Ulva rigida* and *Gracilaria cornea* as dietary ingredients in European seabass (*Dicentrarchus labrax*) juveniles. *Aquaculture*, 252, 85-91.
- Van Alstyne K.L., Wolfe G. V., Freidenburg T. L., Neill A., Hicken C. (2001) Activated defense systems in marine macroalgae: evidence for an ecological role for DMSP cleavage. *Marine Ecology Progress Series Mar Ecol Prog Ser.*, Vol. 213: 53–65.
- Van Tri D. and Thanh Ha N.T., 2004. Impacts of feed on the growth of white leg shrimp larvae *Litopenaeus vannamei* (Boone,1931). Research Institute for Aquaculture No. 3, Fisheries review No. 9/2004.
- Ventura, M.R. & Castañón, J.I.R. (1998) The nutritive value of seaweed (*Ulva lactuca*) for goats. *Small Ruminant Research*, 29, 325-327.
- Viera, M.P., Gómez-Pinchetti, J.L., CourtoisdeVicose, G., Bilbao, A., Suárez, S., Haroun, R.J. & Izquierdo, M.S. (2005) Suitability of three red macroalgae as a feed for the abalone *Haliotis tuberculata coccinea* Reeve. *Aquaculture*, 248(1-4), 75-82.
- Wahbeh, M.I. (1997) Amino acid and fatty acid profiles of four species of macroalgae from Aqaba and their suitability for use in fish diets. *Aquaculture*, 159, 101-109.
- Wong, K.H. & Cheung, P.C.K. (2000) Nutritional evaluation of some subtropical red and green seaweeds. Part I.– Proximate composition, amino acid profiles and some physico-chemical properties. *Food chemistry*, 71, 475-482.
- Wong, K.H. & Cheung, P.C.K. (2001a) Nutritional evaluation of some subtropical red and green seaweeds. Part II. In vitro protein digestibility and amino acid profiles of protein concentrates. *Food chemistry*, 72, 11-17.
- Wong, K. & Cheung, P.Ch. (2001b) Influence of drying treatment on three *Sargassum* species. *Journal of Applied Phycology*, 13(1), 43-50.
- Yeh, S.T., Lee, C.S. & Chen, J.C. (2006) Administration of hot-water extract of brown seaweed *Sargassum duplicatum* via immersion and injection enhances the immune resistance of white shrimp *Litopenaeus vannamei*. *Fish Shellfish Immunol.*, 20(3), 332-345
- Yone, Y., Furuichi, M. & Urano, K. (1986a) Effects of dietary wakame *Undaria pinnatifida* and *Ascophyllum nodosum* supplements on growth, feed efficiency, and proximate compositions of liver and muscle of red sea bream. *Nippon Suisan Gakkaishi*, 52, 1465-1488.
- Yone, Y., Furuichi, M. & Urano, K. (1986b) Effects of wakame *Undaria pinnatifida* and *Ascophyllum nodosum* on absorption of dietary nutrients, and blood sugar and plasma free amino-N levels of red sea bream. *Nippon Suisan Gakkaishi*, 52: 1817-1819.