



Investigación e Innovación en Nutrición Acuícola

**Editores: Lucía Elizabeth Cruz Suárez,
Mireya Tapia Salazar, Martha Guadalupe
Nieto López, David A. Villarreal Cavazos,
Julián Gamboa Delgado, y Carlos A.
Martínez Palacios**

Investigación e Innovación en Nutrición Acuícola

2022, Monterrey, Nuevo León, México

Editores: Lucía Elizabeth Cruz Suárez, Mireya Tapia Salazar, Martha Guadalupe Nieto López, David Alonso Villarreal Cavazos, Julián Gamboa Delgado y Carlos A. Martínez Palacios.

Programa Maricultura
Facultad de Ciencias Biológicas
Universidad Autónoma de Nuevo León 2022

Copias disponibles en:
Universidad Autónoma de Nuevo León
Facultad de Ciencias Biológicas
Programa Maricultura
Cd. Universitaria
San Nicolás de los Garza, Nuevo León
C.P. 66455
Tel.+Fax. 818352 6380
E-mail: lucia.cruzsr@uanl.edu.mx

Para citar alguna parte de ésta obra siga el siguiente estilo:

- Autores del escrito. 2022. Nombre del artículo. Editores: Lucía Elizabeth Cruz Suárez, Mireya Tapia Salazar, Martha Guadalupe Nieto López, David Alonso Villarreal Cavazos, Julián Gamboa Delgado y Carlos A. Martínez Palacios. Investigación e innovación en nutrición acuícola, Universidad Autónoma de Nuevo León, San Nicolás de los Garza, Nuevo León, México, pp.473 ISBN: 978-607-27-1732-9. El cuidado de la presente edición, así como su realización estuvo a cargo de los editores.

La reproducción total o parcial de ésta obra requiere la autorización escrita por los titulares del derecho de autor.

Los editores hacemos extensivo nuestro profundo agradecimiento:

- A las personas que colaboraron en la edición técnica de estas memorias

Directorio

Dr. Santos Guzmán López
Rector

Dr. Juan Paura García
Secretario. General

Dr. Celso José Garza Acuña
Secretario extensión y cultura

Lic. Antonio Ramos Revillas
Director de Editorial Universitaria

Dr. José Ignacio González Rojas
Director de la Facultad de Ciencias Biológicas

Editores

Lucía Elizabeth Cruz Suárez, Denis Ricque Marie, Mireya Tapia Salazar, Martha Guadalupe Nieto López, David Alonso Villarreal Cavazos, Julián Gamboa Delgado, y Carlos A. Martínez Palacios.

Dirección de edición: Programa Maricultura, Facultad de Ciencias Biológicas, Universidad Autónoma de Nuevo León, Dra. Lucía Elizabeth Cruz Suárez, Av. Universidad S/N, Ciudad Universitaria, San Nicolás de los Garza, Nuevo León, C.P. 66455
Email: elicruz@hotmail.com, lucia.cruzsr@uanl.edu.mx
Teléfonos: 52 8183526380

Primera edición 2022. ©Universidad Autónoma de Nuevo León. ©L. Elizabeth Cruz Suárez, ©Mireya Tapia Salazar, ©Martha Guadalupe Nieto López, ©David Alonso Villarreal Cavazos, ©Julián Gamboa Delgado, ©Carlos A. Martínez Palacios.

ISBN:978-607-27-1732-9. El cuidado y edición estuvo a cargo de los editores. El contenido es responsabilidad de los autores.

Párrafo legal: Reservado todos los derechos conforme a la ley. Prohibida la reproducción total o parcial de la obra sin previa autorización por escrito del titular propietario y editor de la obra.

The Role of Fish Nutrition in Improving Human Health and Global Food Security

Albert G.J. Tacon^a & Daniel Lemos^b

^a Aquahana LLC, Honolulu, Hawaii, USA agjtacon@aol.com

^b Universidade de São Paulo, Instituto Oceanográfico, São Paulo, Brazil,

E-mail: dellemos@usp.br

Abstract

Malnutrition represents the world's greatest preventable health challenge, including both undernutrition and overnutrition. The role play by fish and seafood in global food supply and within the Americas is discussed, with aquatic food products offering a much healthier alternative to farmed terrestrial meat products. The paper highlights the important role played by fish nutrition in the formulation of fish and shrimp feeds, and in particular for the nutritional enhancement of farmed fish products through dietary fortification with essential nutrients and improved health attributes,

Keywords: *malnutrition, aquaculture, food supply, human health, nutrient fortification*

The double burden of malnutrition

Malnutrition represents the world's greatest preventable health challenge; the term malnutrition covering both undernutrition (which includes wasting [low weight-for-height], stunting [low height-for-age], underweight [low weight-for age], and vitamin and mineral deficiencies) and overnutrition (which includes overweight, obesity and diet related noncommunicable diseases [NCDs] such as heart disease, hypertension and stroke, diabetes and some cancers).

According to the World Health Organization nearly one in three persons globally suffers from at least one form of malnutrition, with over 1.9 billion adults being overweight or obese (WHO, 2017; 2021a) and 928 million people suffering from severe food insecurity; over half of the world's undernourished living in Asia (418 million) and over a third in Africa (282 million; FAO *et al.*, 2021). Globally, an estimated 155 million children under the age of 5 years are suffer from stunting, while 41 million are overweight, and with around 45% of all deaths among children under 5 years of age being linked to undernutrition (WHO, 2021a).

In contrast to undernutrition, which is mainly linked to low income and poverty, the rapid rise of obesity and diet related NCDs within developed and developing countries has mainly resulted from the increased consumption of lower-cost processed fast foods and high energy-dense diets, coupled with a more sedentary life style (FAO, 2021a; FAO *et al.*, 2021; Tacon *et al.*, 2020; WHO, 2017, 2021a).

Role of fish in global food supply and within the Americas

Although oceans and freshwaters cover more than 71% of our planet, aquatic food products¹ represented less than 3.25 % of total global food supply in 2018 (171.44 million tonnes, live weight equivalent basis), and supplied 108.65 million tonnes of aquatic meat products to the global food basket (FAO, 2021a). Moreover, in terms of nutrient supply, fish and seafood supplied 6.8% of total animal calorific supply, 16.9% of total animal protein supply and 3.2% of total animal fat supply on a global basis in 2018; freshwater fish supplying over 42.9% of total fish and seafood animal protein supply, followed by pelagic fish (20.0%), demersal fish (15.0%), crustaceans (9.1%), other marine fish (5.9%), cephalopods (3.2%), other molluscs (3.2%) and other aquatic animals (0.36%; Table 1, FAO, 2021a).

¹ Includes wild and farmed fish and seafood, including crustaceans, molluscs (including cephalopods), other invertebrate aquatic animals, and aquatic plants;

Table 1. Contribution of fish & aquatic animal products to animal food supply by major geographic region in 2018 (FAO, 2021a)

Region	Total fish food supply 1000 tonnes	Total fish food supply kg/cap/yr	Plant/animal food supply kcal/cap/dy	Food fish supply kcal/cap/dy	Fish protein supply ⁶ g/cap/dy	Fish fat supply g/cap/dy
Asia	108,296 (71.4%)	23.95	2,373/476	40 (8.40%)	6.46 (22.7%)	1.33 (3.79%)
Africa	11,718 (7.7%)	10.27	2,403/200	19 (9.50%)	3.04 (20.0%)	0.70 (5.39%)
Europe	15,997 (10.5%)	21.39	2,436/948	46 (4.85%)	6.48 (11.1%)	1.99 (2.82%)
Oceania	865 (0.57%)	26.75	2,325/981	44 (4.48%)	6.73 (10.4%)	1.60 (2.23%)
Central America	2,111 (1.4%)	12.03	2,409/612	24 (3.92%)	3.66 (9.23%)	0.83 (1.89%)
^a Northern America	8,081 (5.3%)	22.19	2,747/1,013	36 (3.55%)	5.42 (7.55%)	1.29 (1.82%)
^b South America	4,112 (2.7%)	9.71	2,350/734	17 (2.32%)	2.75 (5.79%)	0.62 (1.18%)
^c World	151,641	20.37	2,402/527	36 (6.8%)	5.59 (16.9%)	1.24 (3.25%)

^a Belize, Costa Rica, El Salvador, Guatemala, Honduras, Mexico, Nicaragua, Panama

^b Canada, USA

^c Argentina, Bolivia, Brazil, Chile, Colombia, Ecuador, Guyana, Paraguay, Peru, Suriname, Uruguay, Venezuela

Regionally, the Asian region produced over 71.4% of total global food fish production in 2018, followed by Europe (10.5% by weight), Africa (7.7%), Northern America (5.3%), South America (2.7%), Central America (1.4%), and Oceania, respectively (0.57%; Table 1). However, in contrast to the Asian region where fish and seafood supply is mainly derived from increased aquaculture production (the Asian region producing over 110.03 million tonnes or 91.6% of total global aquaculture production in 2019; FAO, 2021b), fish and seafood supply within the other regions is still largely sourced from wild capture fisheries (FAO, 2020).

In terms of nutrient supply, the highest contribution of fish and seafood products to total nutrient supply was in the Asian region (22.7% of total animal protein supply, 3.79% of total animal fat supply, and 8.40% of total animal calorific supply), followed by the African region (20.0% of total animal protein supply, 5.39% of total animal fat supply, and 9.5% of total calorific supply; Table 1). By contrast, despite a high per capita fish and seafood supply within Oceania (26.75 kg/cap/year), Northern America (22.19 kg/cap/year), and Europe (21.39 kg/cap/year), fish and seafood supplied only 10.4% of total animal protein supply and 2.23% of animal fat supply in Oceania, 7.55% of total animal protein supply and 1.82% of animal fat supply in Northern America, and 11.1% of total animal protein supply and 2.82% of animal fat supply in Europe (Table 1).

By contrast, the lowest contribution of fish and seafood to nutrient supply was in South America (2.32%, 5.79% and 1.18% of total energy, protein and fat supply, respectively), followed by Northern America (3.55%, 7.55% and 1.82% of total energy, protein and fat supply, respectively) and Central America (3.92%, 9.23% and 1.89% of total energy, protein and fat supply, respectively; Table 1). On a country basis, fish and seafood supply in the region varied from a low of 0.9%, 2.1% and 0.5% of total energy, protein and fat supply in Bolivia, to a high of 14.2%, 24.1% and 9.9% of total energy, protein and fat supply in Peru, respectively (Table 2).

Moreover, within most countries where fish and seafood currently play a minor role in animal protein supply (supplying <10% of animal protein supply) there is a general over consumption of energy rich terrestrial animal food products, processed foods, and high energy beverages (sugars and sweeteners supplying over 10% of total calorific supply; Figure 1; Table 2). As a consequence of these dietary food choices it is perhaps not surprising therefore that obesity, heart diseases and diabetes and NCDs have emerged as a major health risk and cause of death within most countries within the region, including Brazil, Mexico and the USA (Tacon *et al.*, 2020; Cameron *et al.*, 2021; Ferreira *et al.*, 2021; Global Obesity Observatory, 2021; Lawrence *et al.*, 2021; Silva *et al.*, 2021; World Bank, 2021b; Fig. 2).




Figure 1. Contribution of the major food groups to total food supply by region (values expressed as % total calorie supply; FAO, 2021a)




Figure 2. Adult obesity 1990-2019 (%) in the United States, Mexico and Brazil. Data source: Global Obesity Observatory (<http://data.worldobesity.org>). Brazil data 2015 and 2019 are author's estimates and also based on Silva *et al.* (2021).

Table 2. Contribution of fish and seafood to food supply within countries within the Americas
(contribution of fish and seafood toward nutrient supply expressed as %; FAO, 2021a)

Country	Total food supply per capita per day						Fish & seafood food supply/capita/day		
	Total calories (kcal)	Animal calories (kcal)	Animal protein (g)	Animal fat (g)	Sugar & sweeteners ¹ (kcal)	Calories	Protein g	Fat g	
Venezuela	2,210	347 (5.8%)	24.57 (12.2%)	24.30 (3.0%)	256 (12.1%)	20	2.99	0.74	
Bolivia	2,412	544 (0.9%)	35.46 (2.1%)	40.97 (0.5%)	308 (12.8%)	5	0.75	0.21	
Guatemala	2,551	289 (2.1%)	21.75 (4.6%)	19.12 (1.1%)	487 (19.1%)	6	1.00	0.21	
Nicaragua	2,582	304 (3.6%)	19.66 (7.8%)	20.59 (2.2%)	407 (15.8%)	11	1.53	0.46	
Ecuador	2,606	535 (2.6%)	30.73 (7.7%)	40.7 (1.2%)	207 (7.9%)	14	2.36	0.49	
Honduras	2,673	344 (1.4%)	21.94 (3.5%)	23.28 (1.0%)	446 (16.7%)	5	0.76	0.24	
El Salvador	2,696	439 (3.4%)	26.91 (8.1%)	30.89 (1.8%)	413 (15.3%)	15	2.19	0.57	
Suriname	2,698	327 (10.1%)	24.71 (19.3%)	22.30 (5.8%)	402 (14.9%)	33	4.78	1.30	
Paraguay	2,768	387 (2.1%)	28.08 (4.4%)	26.16 (1.2%)	286 (10.3%)	8	1.23	0.31	
Peru	2,775	339 (14.2%)	29.78 (24.1%)	19.2 (9.9%)	222 (8.0%)	48	7.18	1.91	
Panama	2,851	619 (5.8%)	42.74 (12.0%)	42.93 (3.5%)	341 (12.0%)	36	5.13	1.49	
Guyana	2,913	470 (8.7%)	35.54 (19.7%)	27.47 (3.8%)	317 (10.9%)	41	7.01	1.05	
Costa Rica	3,029	697 (6.2%)	45.97 (12.5%)	45.53 (4.4%)	503 (16.6%)	43	5.73	2.00	
Chile	3,029	809 (3.0%)	49.6 (7.2%)	62.54 (1.4%)	444 (14.6%)	24	3.56	0.91	
Colombia	3,114	545 (2.6%)	37.31 (5.5%)	37.02 (1.6%)	590 (18.9%)	14	2.05	0.59	
Mexico	3,157	697 (3.9%)	44.78 (9.6%)	50.68 (1.8%)	445 (14.1%)	27	4.30	0.92	
Uruguay	3,202	801 (2.2%)	46.22 (5.7%)	58.48 (1.2%)	477 (14.9%)	18	2.64	0.72	
Brazil	3,301	860 (1.7%)	53.77 (4.5%)	61.44 (0.7%)	409 (12.4%)	15	2.45	0.46	
Argentina	3,307	1,043 (1.2%)	69.05 (3.0%)	76.34 (0.5%)	453 (13.7%)	13	2.08	0.39	
Canada	3,566	888 (4.0%)	56.75 (9.9%)	68.89 (1.8%)	447 (12.5%)	36	5.63	1.22	
USA	3,782	1,027 (3.5%)	73.46 (7.3%)	71.18 (1.8%)	589 (15.6%)	36	5.40	1.30	

¹ Expressed as percent total calories

Fish and seafood – a healthier alternative to terrestrial meat products

Aquatic food products offer a much healthier food source than terrestrial meat products in the global fight against malnutrition and obesity; aquatic animal foods (whether captured or cultured) having a higher protein content on an edible weight basis than most terrestrial meats, a lower caloric density and generally being much leaner than red and processed meats, having the highest content of long-chain omega-3 polyunsaturated fatty acids than any other animal foodstuff, and generally having a higher mineral and vitamin content than most terrestrial meats and processed meat products (Reames 2012; Sargent & Tacon, 1999; Tacon and Metian 2013; USDA, 2018). Moreover, considerable scientific data exists concerning the direct health benefits of consuming aquatic food products, either through the reduced risk of death from coronary heart disease and stroke, the reduced risk of diabetes, the increased duration of gestation and improved visual and cognitive development, the improved neurodevelopment in infants and children when fish is consumed before and during pregnancy, and the reduced risk of thyroid cancer in women through seaweed consumption (FAO/WHO, 2011; Forouhi *et al.*, 2018; He, 2009; Hellberg *et al.*, 2012; Michikawa *et al.*, 2012; Verbeke *et al.*, 2005; Wallin *et al.*, 2012).

Despite the above nutritional and health attributes, there are potential health risks from the consumption of aquatic food products. These can include specific health risks from the consumption of raw/under-cooked fish or shellfish (contaminated with live pathogens or parasites; FAO/WHO, 2011; Hellberg *et al.*, 2012) or from the consumption of aquatic food products sourced from polluted waters (contaminated with environmental pollutants such as heavy metals, persistent organic pollutants [POPs], veterinary drug residues, or micro-plastics; Barber *et al.* 2006; Domingo *et al.* 2007; Lie, 2008; Nicklisch *et al.* 2017; Schmid *et al.* 2007; Tacon & Metain, 2008; Verbeke *et al.* 2005; VKM, 2014; Weber & Goerke, 2003; Xanthos & Walker, 2017) or from the consumption of fish and shellfish sourced from waters containing algal biotoxins (Morabito *et al.* 2018). However, in most instances these risks can be eliminated or greatly reduced by the use of appropriate cooking, handling and/or processing techniques, or through the selection of wild aquatic food products from unpolluted waters. Notwithstanding the above risks, it is generally believed that the higher nutritional value and potential health benefits derived from increased fish consumption far out-way the potential negative risks to human health (FAO/WHO, 2011; VKM, 2014).

Role of fish nutrition in improving human health

It follows from the above discussion that wild-caught fish and seafood products have a greater risk of being contaminated with environmental pollutants and algal biotoxins than farmed aquatic species, where strict controls are usually placed on feed quality and food safety (Barber *et al.*, 2006; Escher *et al.*, 2020; Kwasek *et al.*, 2020; Lie, 2008; Maule *et al.*, 2007; Nicklisch *et al.*, 2017; Schmid *et al.*, 2007; Tacon & Metian, 2008, 2013; USDA, 2018; VKM, 2014; Weber & Goerke, 2003).

In view of the potential presence of environmental pollutants within wild-caught fish and the global stagnation of capture fisheries landings (FAO 2021b), it is clear that aquaculture is the only real long-term sustainable solution for increased fish and seafood production in the short or long term (Tacon, 2020). Moreover, in the case of aquaculture (in contrast to wild capture fisheries), it is also possible to enhance the nutritional profile of the flesh of the cultured by dietary manipulation and/or dietary fortification to meet potential consumer needs (Lie, 2008; Tacon *et al.*, 2020).

Examples of research studies which have successfully demonstrated the nutritional enhancement of farmed fish through dietary fortification are numerous (Figure 2), and examples can be listed as follows:

- Dietary fortification with long chain omega-3 polyunsaturated fatty acids, including Atlantic salmon *Salmo salar* (Kousoulaki *et al.*, 2016; Lie 2008; Nanton *et al.*, 2012; Sprague *et al.*, 2016; Rosenlund *et al.*, 2011; Torstensen *et al.*, 2004), Channel catfish *Ictalurus punctatus* (Manning *et al.*, 2006; Manning *et al.*, 2007), Tilapia *Oreochromis* sp. (Jiratanan 2007; Jorge *et al.*, 2021; Ng *et al.*, 2013; Petenuci *et al.*, 2018; Sarker *et al.*, 2016; Stoneham *et al.*, 2018; Watters *et al.*, 2013) and shrimp *Litopenaeus vannamei* (Kumar *et al.*, 2018; Nonwachai *et al.*, 2010; Wang *et al.*, 2015);
- Dietary fortification with fat soluble vitamins, including vitamin E: Atlantic salmon *S. salar* (Harare *et al.*, 1998; Sigurgisladottir *et al.*, 1994; Waagbø *et al.*, 1993), Rainbow trout *O. mykiss* (Mihalca *et al.*, 2011);
- Dietary fortification with essential trace minerals, including 1) Selenium: African catfish *Clarias gariepinus* (Luten *et al.*, 2008; Schram *et al.*, 2008), Atlantic salmon *S. salar* (Berntssen *et al.*, 2018; Lorentzen *et al.*, 1994; Sele *et al.*, 2018), common carp and gilthead seabream (Barbosa *et al.*, 2020), Rainbow trout *O. mykiss* (Ribeiro *et al.*, 2017); 2) Iodine, including Atlantic salmon *S. salar* (Julshamn *et al.*, 2006; Schmid *et al.*, 2003), Chars *Salvelinus* sp. (Schmid *et al.*, 2003),

common carp and gilthead seabream (Barbosa *et al.*, 2020), Rainbow trout *Oncorhynchus mykiss* (Ribeiro *et al.*, 2017); and 3) trivalent Chromium: Nile Tilapia *O. niloticus* (Li *et al.*, 2018); and

- Dietary fortification with essential and non-essential amino acids, including Taurine: African catfish *Clarias gariepinus* (Luten *et al.*, 2008) and Rainbow trout *O. mykiss* (Anderson 1992; Aksnes *et al.*, 2006).




Figure 2. Tissue levels in farmed fish which can be manipulated through dietary fortification

Moreover, in addition to the above, the concentration of environmental pollutants, including heavy metals and POPs, can be reduced within farmed fish (if deemed a problem) through the dietary substitution of contaminated fish oils and fish meals with less contaminated lipid and protein meals (Bell *et al.*, 2005; Berntssen *et al.*, 2005; Berntssen *et al.*, 2011; Haldorsen *et al.*, 2017; Heshmati *et al.*, 2019; Turchini *et al.*, 2019).

Need for improved communication & consumption of farmed aquatic food products

Fish and seafood needs to be viewed by policy makers and governments as an important contributor to the global food basket, and the increased consumption of fish and seafood products actively promoted as a high-health and heart-healthy alternative to processed foods and terrestrial animal food products; the latter being particularly true for those countries suffering from a high incidence of obesity, coronary heart disease, diabetes and associated ailments (Henry *et al.*, 2020; HHP, 2011; Mozaffarian and Ludwig 2010; NHMRC, 2013; Pan *et al.*, 2012; Skerrett and Willett 2010; Tacon, Lemos & Metian, 2020; USDHHS/USDA, 2015; Wallin *et al.*, 2012, 2016; Willett *et al.*, 2019).

Table 3 presents recommended targets for a 2,500 Kcal healthy reference diet, including the possible ranges for different food groups, including fish. Moreover, it has been estimated that dietary changes from current high energy dense diets toward more healthy diets are likely to have significant health benefits, including averting between 7.4 to 10.8 million premature deaths per year (Willett *et al.*, 2019).

Table 3. Recommended targets for a healthy reference diet, with possible ranges, for an intake of 2,500 Kcal/day (compiled from Willet *et al.*, 2019)

Food group	Food subgroup	Reference diet (g/day)	Kcal/day	Possible ranges (g/day)
Whole grains	Rice, wheat, corn, beans, lentils	232	811	Total grains 0 to 60% of energy
Starchy roots	Potatoes, cassava	50	39	0 to 100
Vegetables	All vegetables	300		200 to 600
Fruits	All fruit	200	126	100 to 300
Dairy foods	Dairy foods	250	153	0 to 500
Protein sources	Beef, lamb	7	15	0 to 14
	Pork	7	15	0 to 14
	Chicken, other poultry	29	62	0 to 58
	Eggs	13	19	0 to 25
	Fish	28	40	0 to 100
	Beans, lentils & peas	50	172	0 to 100
	Soy foods	25	112	0 to 50
	Tree nuts & peanuts	50	291	0 to 75
Added fats	Unsaturated plant oils	40	354	20 to 80
	Dairy fats	0	0	
	Lard/tallow	5	36	0 to 5
Added sugars	All sweeteners	31	120	0 to 31

References cited

- Aksnes A, Hope B, Albrektsen S. (2006). Size-fractionated fish hydrolysate as feed ingredient for rainbow trout (*Oncorhynchus mykiss*) fed high plant protein diets. II: Flesh quality, absorption, retention and fillet levels of taurine and anserine. *Aquaculture*. 261(1):318–326. doi: 10.1016/j.aquaculture.2006.07.026
- Anderson RA. (1992). Chromium, glucose tolerance, and diabetes. *Biol Trace Elem Res*. 32(1-3):19–24. doi:10.1007/BF02784583
- Barber, L.B., S.H. Keefe., R.C. Antweiler., H.E. Taylor & R.D. Wass. (2006). Accumulation of contaminants in fish from waste- water treatment. *Environ Sci Technol*. 40(2):603–611. doi: 10.1021/es0514287
- Barbosa, V., Maulvault, A.L., Anacleto, P., Santos, M., Mai, M., Oliveira, H., Delgado, I., Coelho, I., Barata, M., Araújo-Luna, R., Ribeiro, L., Eljasik, P., Sobczak, M., Sadowski, J., Tórz, A., Panicz, R., Dias, J., Pousão-Ferreira, P., Carvalho, M.L., Martins, M., Marques, A. (2020). Enriched feeds with iodine and selenium from natural and sustainable sources to modulate farmed gilthead seabream (*Sparus aurata*) and common carp (*Cyprinus carpio*) fillets elemental nutritional value. *Food and chemical toxicology*, 140: <https://doi.org/10.1016/j.fct.2020.111330>
- Bell JG, McGhee F, Dick JR, Tocher DR. (2005). Dioxin and dioxin-like polychlorinated biphenyls (PCBs) in Scottish farmed salmon (*Salmo salar*): effects of replacement of dietary marine fish oil with vegetable oils. *Aquaculture*. 243(1-4):305–314. doi:10.1016/j.aquaculture.2004.10.016
- Berntssen MHG, Lundebye A-K, Torstensen BE. (2005). Reducing the levels of dioxins and dioxin-like PCBs in farmed Atlantic salmon by substitution of fish oil with vegetable oil in the feed. *Aquac Nutr*. 11(3):219–231. doi: 10.1111/j.1365-2095.2005.00345.x
- Berntssen MHG, Maage A, Julshamn K, Oeye BE, Lundebye A-K. (2011). Carry-over of dietary organochlorine pesticides, PCDD/Fs, PCBs, and brominated flame retardants to Atlantic salmon (*Salmo salar* L.) fillets. *Chemosphere*. 83(2):95–103. doi:10.1016/j.chemosphere.2011.01.017
- Berntssen M, Betancor MB, Caballero MJ, Hillestad M, Rasinger JD, Hamre K, Sele V, Amlund H, Ørnsrud R. (2018). Safe limits of selenomethionine and selenite supplementation to plant-based Atlantic salmon feeds. *Aquaculture*. 495:617–630. doi:10.1016/j.aquaculture.2018. 06.041
- Cameron, N.A., Petito, L.C., McCabe, M., Allen, N.B., O'Brien, M.J., Carnethon, M.R., Khan, S.S. (2021). Quantifying the sex-race/ethnicity-specific burden of obesity on incident diabetes mellitus in the United States, 2001 to 2016: MESA and NHANES. *J. Am. Heart Assoc*. 2021; 10:e018799. DOI: 10.1161/JAHA.120.018799
- Dey, M.M., Garcia, Y.T., Praduman, K., Piumsombun, S., Haque, M.S., Li, L., Radam, A., Senaratne, A., Khiem, N.T. and Koeshendrajana, S. (2008). Demand for fish in Asia: a cross-country analysis. *Australian Journal of Agricultural and Resource Economics*, 52(3), pp.321-338.
- Domingo, J.L., A. Bocio., G. Falco & J.M. Llobet. (2007). Benefits and risks of fish consumption. Part 1. A quantitative ana- lysis of the intake of omega-3 fatty acids and chemical contaminants. *Toxicology*. 230(2-3):219–226. doi:10.1016/ j.tox.2006.11.054

- Escher, B.I., H.M. Stapleton & E.L. Schymanski. (2020). Tracking complex mixtures of chemicals in our changing environment. *Science*. 367(6476):388–392. doi:10.1126/science. aay6636
- FAO. (2020). The State of World Fisheries and Aquaculture 2020. Sustainability in action. Rome. <https://doi.org/10.4060/ca9229en>
- FAO. (2021a). FAO Food Balance Sheet. <http://www.fao.org/faostat/en/#data/FBS>. Last update February 19, 2020.
- FAO. (2021b). FishStatJ, a tool for fishery statistics analysis. Release: 4.01.1 May 2021. Universal Software for Fishery Statistical Time Series. Global aquaculture production: Quantity 1950–2019; Value 1950–2019; Global capture production. Rome, Italy: FAO. p. 1950–2019.
- FAO/WHO. (2011). Report of the joint FAO/WHO expert consultation on the risks and benefits of fish consumption, 25–29 January 2010. Rome, Italy: FAO. p. 50. Fisheries and Aquaculture Report No. 978 FIPM/R978 (En).
- FAO, IFAD, UNICEF, WFP & WHO. (2021). The State of Food Security and Nutrition in the World 2021. Transforming food systems for food security, improved nutrition and affordable healthy diets for all. Rome, FAO. <https://doi.org/10.4060/cb4474en>
- Ferreira, C.M. Reis, N.D., Castro, A.O., Höfelmann, D.A., Kodaira, K., Silva, M.T., Galvao, T.F. (2021). Prevalence of childhood obesity in Brazil: systematic review and meta-analysis. *Jornal de Pediatria*, <https://doi.org/10.1016/j.jpmed.2020.12.003>
- Forouhi, N.G., A. Misra., V. Mohan., R. Taylor & W. Yancy. (2018). Dietary and nutritional approaches for prevention and management of type 2 diabetes. *BMJ*. 361:k2234.
- Global Obesity Observatory (2021). https://data.worldobesity.org/country/mexico-139/#data_trends
- Kumar, V Habte-Tsion. H.M., Allen, K.M., Bowman, B.A., Thompson, K.R., El-Haroun, E., Filer, K., Tidwell, J.H. (2018). Replacement of fish oil with *Schizochytrium* meal and its impacts on the growth and lipid metabolism of Pacific white shrimp (*Litopenaeus vannamei*). *Aquaculture Nutrition*, 24: 1769-1781.
- Haldorsen A-K, Lock E-J, Rasinger JD, Nøstbakken OJ, Hannisdal R, Karlsbakk E, Wennevik V, Madhun AS, Madsen L, Graff IE, *et al.* (2017). Lower levels of Persistent Organic Pollutants, metals and the marine omega 3-fatty acid DHA in farmed compared to wild Atlantic salmon (*Salmo salar*). *Environ Res*. 155:49–59. doi:10.1016/j.envres.2017.01.026
- Harare K, Berge RK, Lie Ø. (1998). Oxidative stability of Atlantic salmon (*Salmo salar*, L.) fillet enriched in alpha -, gamma -, and delta -tocopherol through dietary supplementation. *Food Chem*. 62(2):173–178. doi:10.1016/S0308-8146(97)00209-4
- He, K. (2009). Fish, long-chain omega-3 polyunsaturated fatty acids and prevention of cardiovascular disease—eat fish or take fish oil supplement? *Prog Cardiovasc Dis*. 52(2): 95–114. doi:10.1016/j.pcad.2009.06.003
- Hellberg, R.S., C.A.M. De Witt & M.T. Morrissey. (2012). Risk-benefit analysis of seafood consumption: a review. *Compr Rev Food Sci F*. 11(5):490–517. doi:10.1111/j.1541-4337. 2012.00200.x
- Henry CJ, Kaur B, Quek RYC. (2020). Are Asian foods as “fattening” as western-styled fast foods? *European J Clin Nutr*. 74(2):348–350. 2010). doi:10.1038/s41430-019- 0537-3
- Heshmati A, Sadati R, Ghavami M, Khaneghah AM. (2019). The concentration of potentially toxic elements (PTEs) in muscle tissue of farmed Iranian rainbow trout (*Oncorhynchus mykiss*), feed, and water samples collected

- from the West of Iran: A risk assessment study. *Environ Sci Pollut Res.* 26(33):34584–34593. doi:10.1007/s11356-019-06593-x
- Harvard Health Publishing (HHP). (2011) September 14. Harvard to USDA: check out the healthy eating plate. Harvard Medical School. [accessed 2020 February 28]. <https://www.health.harvard.edu/blog/harvard-to-usda-check-out-the-healthy-eating-plate-201109143344>.
- Jiratanan T. (2007). Development of omega-3 fatty acid enhanced tilapia (*Oreochromis niloticus*) (alpha-Linolenic acid). *Dissertation Abstr Int.* 68(3, suppl. B):92.
- Jorge, T.B.F., Moura, G.S., Ribeiro-Junior, V., Donzele, J.L., Pedreira, M.M.. 2021, Sousa, T.V., Lanna, E.A.T. 2021. Effects of dietary supplementation time with *Schizochytrium* microalgae meal on growth, meat quality and fatty acid composition of Nile tilapia. *Aquaculture Research*, doi.org/10.1111/are.15597
- Julshamn K, Maage A, Waagbo R, Lundebye A-K. (2006). A preliminary study on tailoring of fillet iodine concentrations in adult Atlantic salmon (*Salmo salar* L.) through dietary supplementation. *Aquac Nutrition.* 12(1):45–51. doi:10.1111/j.1365-2095.2006.00380.x
- Kousoulaki K, Mørkøre T, Nengas I, Berge RK, Sweetman J. (2016). Microalgae and organic minerals enhance lipid retention efficiency and fillet quality in Atlantic salmon (*Salmo salar* L. *Aquaculture.* 451:47–57. doi:10.1016/j.aquaculture.2015.08.027
- Kwasek, K., A.L. Thorne-Lyman & M. Phillips. (2020). Can human nutrition be improved through better fish feeding practices? A review paper. *Crit Rev Food Sci Nutr.* 25:1–14.
- Lawrence, J.M., Divers, J., Isom, S., Saydah, S., Imperatore, G., Pihoker, K., Marcovina, S.M., Mayer-Davis, E.J., Hamman, R.F., Dolan, L., Dabelea, D., Pettitt, D.J., Liese, A.D. (2021). Trends in prevalence of type 1 and type 2 diabetes in children and adolescents in the US, 2001-2017. *JAMA* 326(8):717-727.
- LiH, MengX, WanW, LiuH, SunM, WangH, WangJ. (2018). Effects of chromium picolinate supplementation on growth, body composition, and biochemical parameters in Nile tilapia *Oreochromis niloticus*. *Fish Physiol Biochem.* 44(5):1265–1274. doi:10.1007/s10695-018-0514-0
- Lie, Ø. editor. (2008). *Improving farmed fish quality and safety.* Cambridge: Woodland Publishing Ltd. p. 648.
- Lorentzen M, Maage A, Julshamn K. (1994). Effects of dietary selenite or selenomethionine on tissue selenium concentrations of Atlantic salmon (*Salmo salar*). *Aquaculture.* 121(4):359–367. doi:10.1016/0044-8486(94)90270-4
- Luten J, Schram E, and E. Elvevoll (2008). Tailor-made functional seafood for consumers: dietary modulation of selenium and taurine in farmed fish pp.343–362. In: Lie, Ø. editor. *Improving farmed fish quality and safety.* Cambridge: Woodland Publishing Ltd. p. 648.
- Manning BB, Li MH, Robinson EH. (2007). Feeding Channel catfish, *Ictalurus punctatus*, diets amended with refined marine fish oil elevates omega-3 highly unsaturated fatty acids in fillets. *J World Aquaculture Soc.* 38(1):49–58. doi:10.1111/j.1749-7345.2006.00072.x
- Manning BB, Li MH, Robinson EH, Peterson BC. (2006). Enrichment of channel catfish (*Ictalurus punctatus*) fillets with conjugated linoleic acid and omega-3 fatty acids by dietary manipulation. *Aquaculture.* 261(1):337–342. doi: 10.1016/j.aquaculture.2006.07.029

- Maule, A.G., A.L. Gannam & J.W. Davis. (2007). Chemical contaminants in fish feeds used in federal salmonid hatcheries in the USA. *Chemosphere*. 67(7):1308–1315. doi:10.1016/j.chemosphere.2006.11.029
- Mihalca GL, Tit O, Tit M, Mihalca A. (2011). Effect of dietary alpha-tocopheryl acetate on alpha-tocopherol content of novel omega-3-enhanced farmed rainbow trout (*Oncorhynchus mykiss*) fillets. *J Agroalim Processes Technol*. 17(3):295–302.
- Michikawa T, Inoue M, Shimazu T, Sawada N, Iwasaki M, Sasazuki S, Yamaji T, Tsugane S. (2012). Seaweed consumption and the risk of thyroid cancer in women: The Japan Public Health Center-based Prospective Study. *Eur J Cancer Prev*. 21(3):254–260. doi:10.1097/CEJ.0b013e32834a8042
- Morabito, S., Silvestro, S. and Faggio, C. (2018). How the marine biotoxins affect human health. *Natural Product Research*, 32(6):621-631.
- Mozaffarian D, Ludwig DS. (2010). Dietary guidelines in the 21st century: a time for food. *JAMA*. 304(6):681–682. doi:10.1001/jama.2010.1116
- Nanton DA, Ruohonen K, Robb DHF, El-Mowafi A, Hartnell GF. (2012). Effect of soy oil containing stearidonic acid on growth performance and fillet fatty acid composition of Atlantic salmon. *Aquacult Nutr*. 18(6):640–650. doi:10.1111/j.1365-2095.2011.00922.x
- Ng W-K, Chong C-Y, Wang Y, Romano N. (2013). Effects of dietary fish and vegetable oils on the growth, tissue fatty acid composition, oxidative stability and vitamin E content of red hybrid tilapia and efficacy of using fish oil finishing diets. *Aquaculture*. 372-375:97–110. doi:10.1016/j.aquaculture.2012.10.030
- National Health and Medical Research Council (NHMRC). (2013). *Eat for health: Australian dietary guidelines*. Canberra, Australia: National Health and Medical Research Council. p. 210.
- Nicklisch, S.C.T., L.T. Bonito., S. Sandin & A. Hamdoun. (2017). Geographic differences in persistent organic pollutant levels of Yellowfin Tuna. *Environ Health Perspect*. 125(6): 067014. doi:10.1289/EHP518
- Nonwachai T, Purivirojkul W, Limsuwan C, Chuchird N, Velasco M, Dhar AK. (2010). Growth, nonspecific immune characteristics, and survival upon challenge with *Vibrio harveyi* in Pacific white shrimp (*Litopenaeus vannamei*) raised on diets containing algal meal. *Fish Shellfish Immun*. 29(2):298–304. doi:10.1016/j.fsi.2010.04.009
- Pan A, Malik V, Hu FB. (2012). Exporting diabetes to Asia: the impact of western-style fast food. *Circulation*. 126(2): 163–165. doi:10.1161/CIRCULATIONAHA.112.115923
- Petenuci ME, Schneider VVA, Lopes AP, Gonçalves RM, Dos Santos VJ, Matsushita M, Visentainer JV. (2018). Effect of alpha-linolenic acid sources in diets for Nile Tilapia on fatty acid composition of fish fillet using principal component analysis. *J Aquat Food Prod Technol*. 27(4):464–476. doi:10.1080/10498850.2018.1448917
- Reames E. (2012). *Nutritional benefits of seafood*. Stoneville, MS: Southern Regional Aquaculture Center (SRAC) Publication No. 7300. p. 6.
- Ribeiro A.R., Goncalves A, Bandarra N, Nunes ML, Dinis MT, Dias J, Rema P. (2017). Natural fortification of trout with dietary macroalgae and selenised-yeast increases the nutritional contribution in iodine and selenium. *Food Res Internat*. 99(3):1103–1109. doi:10.1016/j.foodres.2016. 10.030

- Rosenlund, G., G. Corraze, M. Izquierdo, and B.E. Tortensen. (2011). The effects of fish oil replacement on nutritional and organoleptic qualities of farmed fish, pp.487-522. In: Turchini G.M., Ng W.-K., Tocher D.R., editors. Fish oil replacement and alternative lipid sources in aquaculture feeds. CRC Press, Taylor & Francis Group. p. 551.
- Sargent, J.R. & A.G.J. Tacon. (1999). Development of farmed fish: a nutritionally necessary alternative to meat. Proc Nutr Soc. 58(2):377–383. doi:10.1017/S0029665199001366
- Sarker PK, Kapuscinski AR, Lanois AJ, Livesey ED, Bernhard KP, Coley ML. (2016). Towards sustainable aqua-feeds: complete substitution of fish oil with marine microalga *Schizochytrium* sp. improves growth and fatty acid deposition in juvenile Nile Tilapia (*Oreochromis niloticus*). Plos One. 11(6):e0156684. doi:10.1371/journal.pone.0156684
- Schmid S, Ranz D, He ML, Bukard S, Lukowics MV, Reiter R, Arnold R, LeDeit H, David M, Rambeck WA. (2003). Marine algae as natural source of iodine in the feeding of freshwater fish: a new possibility to improve iodine supply in man. Rev Med Vet. 154:645–648.
- Schmid, P., M. Kohler., E. Gujer., M. Zennegg & M. Lanfranchi. (2007). Persistent organic pollutants, brominated flame retardants and synthetic musks in fish from remote alpine lakes in Switzerland. Chemosphere. 67(9):S16–S21. doi:10.1016/j.chemosphere.2006.05.080
- Schram E, Pedrero Z, Camara C, van der Heul JW, Luten BJ. (2008). Enrichment of African catfish with functional selenium originating from garlic. Aquaculture Res. 39(8): 850–860. doi:10.1111/j.1365-2109.2008.01938.x
- Sele V, Ørnsrud R, Sloth JJ, Berntssen MHG, Amlund H. (2018). Selenium and selenium species in feeds and muscle tissue of Atlantic salmon. J Trace Elem Med Bio. 47: 124–133. doi:10.1016/j.jtemb.2018.02.005
- Sigurgisladottir S, Parrish CC, Lall SP, Ackman RG. (1994). Effects of feeding natural tocopherols and astaxanthin on Atlantic salmon (*Salmo salar*) fillet quality. Food Res Internat. 27(1):23–32. doi:10.1016/0963-9969(94)90174-0
- Silva, L.E.S., Oliveira, M.M., Stopa, S.R., Gouvea, E.C.D.P., Ferreira, K.R.D., Santos, R.O., Valença Neto, P.F., Macário, E.M., Sardinha, L.M.V. (2021). Temporal trend of overweight and obesity prevalence among Brazilian adults, according to sociodemographic characteristics, 2006-2019. Epidemiol. Serv. Saude, Brasília, 30(1):e2020294, 2021. Doi: 10.1590/S1679-49742021000100008
- Skerrett PJ, Willett WC. (2010). Essentials of healthy eating: a guide. J Midwifery Womens Health. 55(6):492–501. doi: 10.1016/j.jmwh.2010.06.019
- Sprague M, Dick JR, Tocher DR. (2016). Impact of sustainable feeds on omega-3 long-chain fatty acid levels in farmed Atlantic Salmon, 2006–2015. Sci Rep. 6(1):21892. doi:10.1038/srep21892
- Stoneham TR, Kuhn DD, Taylor DP, Neilson AP, Smith SA, Gatlin DM, Chu H, O’Keefe SF. (2018). Production of omega-3 enriched tilapia through the dietary use of algae meal or fish oil: Improved nutrient value of fillet and offal. Plos One. 13(4):e0194241. doi:10.1371/journal.pone.0194241
- Tacon, A.G.J. (2020). Trends in global aquaculture and aqua-feed production: 2000–2017. Rev Fish Sci Aquacult. 28(1):43–56. doi:10.1080/23308249.2019.1649634
- Tacon, A.G.J. & M. Metian. (2008). Aquaculture feed and food safety: the role of FAO and Codex Alimentarius. Ann Newyork Acad Sci. 1140(1):50–59. doi:10.1196/annals.1454.003

- Tacon A.G.J. & M. Metian. (2013). Fish Matters: importance of aquatic foods in human nutrition and global food supply. *Rev Fish Sci.* 21(1):1–17.
- Tacon, A.G.J. & M. Metian. (2017). Food matters: fish, income, and food supply - a comparative analysis, *Reviews in Fisheries Science & Aquaculture*, 26(1):15-28.
- Tacon, A.G.J., D. Lemos, and M. Metain. (2020). Fish for health: improved nutritional quality of cultured fish for human consumption. *Reviews in Fisheries Science & Aquaculture*, 28(4):449–458.
- Torstensen B, Froyland L, Ormsrud R, Lie O. (2004). Tailoring of a cardioprotective fillet fatty acid composition of Atlantic salmon (*Salmo salar*) fed vegetable oils. *Food Chem.*87(4):567–580. doi:10.1016/j.foodchem.2004.01.009
- Turchini GM, Trushenski JT, Glencross BD. (2019). Thoughts for the future of aquaculture nutrition: realigning perspectives to reflect contemporary issues related to judicious use of marine resources in aquafeeds. *North Am J Aquacult.* 81(1):13–39. doi:10.1002/naaq.10067
- US Department of Agriculture (USDA). (2018). Composition of Foods: raw, processed, prepared. USDA National Nutrient Database for Standard Reference, Legacy April 2018, Slightly revised July 2018. <http://www.ars.usda.gov/nutrientdata>.
- U.S. Department of Health & Human Services and U.S. Department of Agriculture (USDHHS/USDA). (2015). 2015–2020 Dietary Guidelines for Americans. 8th Edition. December 2015. <http://health.gov/dietaryguidelines/2015/guidelines/>.
- Verbeke, W., I. Sioen., Z. Pieniak., J. Van Camp & S. De Henauw. (2005). Consumer perception versus scientific evidence about health benefits and safety risks from fish consumption. *Public Health Nutr.*8(4):422–429. doi:10.1079/PHN2004697
- Vitenskapskomiteen (VKM). (2014). Benefit-risk assessment of fish and fish products in the Norwegian diet – an update. Scientific Opinion of the Scientific Steering Committee. VKM Report 15. Oslo, Norway. p. 293. ISBN: 978-82-8259-159-1. www.vkm.no.
- Waagbø R, Sandnes K, Torrissen OJ, Sandvin A, Lie Ø. (1993). Chemical and sensory evaluation of fillets from Atlantic salmon (*Salmo salar*) fed three levels of N-3 polyunsaturated fatty acids at two levels of vitamin E. *Food Chem.*46(4):361–366. doi:10.1016/0308-8146(93) 90005-Z
- Wallin, A., D. Di Giuseppe., N. Orsini., P.S. Patel., N.G. Forouhi & A. Wolk. (2012). Fish consumption, dietary long-chain n-3 fatty acids, and risk of type 2 diabetes-systemic review and met-analysis of prospective studies. *Diabetes Care.*35(4):918–929. doi:10.2337/dc11-1631
- Wang, Y., Li, M., Filer, K., Xue, Y., Ai, Q., Mai, K. (2016). Replacement of fish oil with a DHA-rich *Schizochytrium* meal on growth performance, activities of digestive enzyme and fatty acid profile of Pacific white shrimp (*Litopenaeus vannamei*) larvae. *Aquaculture Nutrition*, 23:1113–1120.
- Watters CA, Rosner LS, Franke AA, Dominy WD, Klinger- Bowen R-E, Tamaru CS. (2013). Watters CA, Rosner LS, Franke AA, Dominy WD, Klinger- Bowen R-E, Tamaru CS. 2013. Nutritional Enhancement of long-chain omega-3 fatty acids in Tilapia (*Oreochromis honorum*). *Bamidgeh.* 65(869):1–7.
- Weber, K. & H. Goerke. (2003). Persistent organic pollutants (POPs) in Antarctic fish: levels, patterns, changes. *Chemosphere.* 53(6):667–678. doi:10.1016/S0045-6535(03) 00551-4

- WHO. (2017). The double burden of malnutrition. Policy brief. Geneva: World Health Organization; 2017. WHO/NMH/NHD/17.3, 11p.
- WHO. (2021a). Malnutrition, 9 June 2021. <https://www.who.int/news-room/fact-sheets/detail/malnutrition>
- Willett, W., Rockström, J., Loken, B., Springmann, M., Lang, T., Vermeulen, S., Garnett, T., Tilman, D., DeClerck, F., Wood, A. and Jonell, M., et al. (2019). Food in the Anthropocene: the EAT–Lancet Commission on healthy diets from sustainable food systems. *The Lancet*, 393(10170), pp.447-492.
- World Bank (2021b). Percentage adults living with obesity, selected countries, 1960-2019. https://data.worldobesity.org/country/mexico-139/#data_trends
- Xanthos, D. & T.R. Walker. (2017). International policies to reduce plastic marine pollution from single-use plastics (plastic bags and microbeads): A review. *Mar Pollut Bull.* 118(1-2):17–26. doi:10.1016/j.marpolbul.2017.02.048